

INDEX

1–9

1- $\bar{3}$ -3- $\bar{1}$ binomial sequence, 224–226, 227f
 180° (π) pulse, 16, 17f. *See also*
 Adiabatic pulse
 Bloch-Siegert shift for, 171
 catenation of, 607, 641
 composite, 174–179, 178f, 203–205
 decoupling by, 94–96, 203–204
 excitation null for, 170
 excitation profile for, 167f, 178f
 EXORCYCLE for, 266, 309, 313
 Gaussian, 182f, 184f
 imperfect, 176, 178f, 309, 547
 inversion by, 16, 17f, 167f, 169–170
 off-resonance effects for, 167f, 169, 178f
 nonresonant phase shift for, 171
 pulsed field gradients and, 313–314
 R-SNOB, 182f, 184f, 186
 RE-BURP, 182f, 184f, 186
 refocusing by, 93–94, 265–266, 703
 selective, 181, 182f, 184f, 186
 sinc, 182f, 184f
 spin-echo and, 93–94

1D (One-dimensional) ^1H NMR,
 263–267, 406–409
 excitation sculpting, 229f, 230, 233f
 inversion-recovery, 258–259, 260f
 jump-return, 224–225, 227f, 233f, 261f
 one-pulse acquire, 263, 265, 267f
 peptide, of, 272f
 spin-echo, Hahn, 259, 228f, 229f, 261f, 265–267, 267f, 407f, 783f
 ubiquitin, of, 150f, 233f, 235f, 260f, 261f, 267f, 272f, 407f, 783f
 ubiquitin, denatured, of, 407f
 water-flipback, 229f, 231–232
 WATERGATE, 229f, 230
 2D (Two-dimensional) NMR, 273–280.
 See also individual NMR experiments
 3D NMR, compared with, 327–229, 525–526
 2D spectrum, schematic, 277
 2Q. *See* Double-quantum coherence
 2Q (Double-quantum) NMR, 465–481
 cosy, comparison with, 475–476
 cross-peak fine structure in, 466–471
 direct peaks in, 469
 experimental protocol for, 473–474

- 2Q (Double-quantum) NMR
(Continued)
 $^1\text{H}^\alpha$ - $^1\text{H}^\beta$ fingerprint region in, 478, 479f
 lineshapes for, 467–471, 470f, 471f, 472t
 processing of, 474
 product operator analysis for, 466–473
 pulse sequence for, 465f
 remote peaks in, 469, 476–477, 478f
 three-spin system, of, 467–468
 two-spin system, of, 466–467, 467f
 ubiquitin, of, 467f, 477f, 478f, 479f, 480f
- 2QF-COSY (Double-quantum filtered COSY), 437–449
 aliasing and folding in, 444
 artifact suppression in, 444–445, 445f
 cross-peak fine structure in, 441–444, 456f
 experimental protocol for, 444–445
 fingerprint regions in, 418–419, 419f, 446
 lineshapes in, 440, 444
 processing of, 445–446
 product operator analysis for, 440–440
 pulse sequence for, 437f, 439f
 S/N ratio and, 447–448, 448f
 three-spin system, of, 442–444
 two-spin system, of, 440–442
 ubiquitin, of, 445f, 446f, 449f, 450f, 453f, 454f, 462f, 480f
- 3D (Three-dimensional) NMR, 525, 581.
See also individual NMR experiments
 2D NMR, compared with, 327–329, 525–526
 4D NMR, compared with, 330, 655
 deuteration and, 733–734
 3D spectrum, schematic, 328f, 329f, 586f
- 3Q. *See* Three-quantum coherence
- 3Q (Three-quantum) NMR, 481–486
 experimental protocol for, 483
 product operator analysis for, 481–483
 processing of, 483
 pulse sequence for, 465f
 ubiquitin, of, 484f
- 3QF-COSY, 449–455
 cross-peak fine structure in, 456f
 experimental protocol for, 451–452
 fingerprint region in
 $^1\text{H}^\alpha$ - $^1\text{H}^\beta$ fingerprint region, 453–455, 454f, 455f
 methyl, 453f
 forbidden cross-peaks, 453
 product operator analysis for, 449–451
 processing of, 452
 pulse sequence for, 437f
 ubiquitin, of, 453f, 454f, 455f, 462f
- 4D NMR. *See also individual NMR experiments*
 3D NMR, compared with, 330, 655
 deuteration and, 733–734
 4D spectrum, schematic, 330f
 90° ($\pi/2$) pulse, 16, 17f, 19f
 excitation null for, 169–170
 off-resonance effects for, 166–169, 167f, 168f
 selective, 183f

A

- AB spin system, 26, 62–64. *See also* Two-spin system
- Absorptive antiphase lineshape, 413f, 421, 423f, 424–426, 424f, 425f, 470f
 COSY and, 411–412
 representation in spectra, 406
- Absorptive lineshape, 18, 139–140, 140f, 317, 320f
- Acquisition period, 17, 19f, 56, 124, 271, 275f, 279, 280. *See also* Indirect evolution period
 initial delay for, 266
 maximum, 144–145, 158
- Active nuclei, 2, 3t
- Active scalar coupling, 91, 282–283, 412
- ADC (Analog-to-digital converter), 115f, 123, 127–129, 321
- Adiabatic condition, 192
- Adiabatic following, 189–190
- Adiabatic full-passage (AFP), 199
- Adiabatic half-passage (AHP), 199
- Adiabatic pulse, 189–201, 709f

- CHIRP, 197
- dephasing ZQ coherence by, 314
- half-filter, in, 768–769
- hyperbolic secant, 198
- magnetization trajectories during, 195f
- sech/tanh, 198
- WURST, 198–199, 200f
- Adiabatic isotropic mixing, 491–493, 492f
- Adiabatic rapid passage, 189–190
- Adiabatic relaxation, 336–337, 349–350
- Adiabatic spin decoupling, 209–212, 213f, 216f, 758
- Adjoint operator, 32
- AFP (Adiabatic full passage), 199
- Aggregation, 19, 236, 408
- AHP (Adiabatic half-passage), 199
- Aliasing and folding, 125–126, 323–326, 325f, 550f, 551f
- 2QF-COSY, in, 444
- ^{13}C spectrum, of, 551f, 769
- heteronuclear correlation and, 549–552
- multidimensional NMR, in, 323–326
- ^{15}N spectrum, of, 551f
- Allowed transition, 25, 25f, 77f
- Amide proton exchange
 - backbone correlations missing due to, 418
 - exchange cross-peaks from, 512–513
 - hydrogen bond restraints from, 805–806
 - intrinsic rates of, 223–224, 224f
 - line broadening by, 406
 - measuring, 499–500
 - perdeuteration and, 727–728
 - protein-ligand interface and, 760
 - saturation transfer by, 223, 409, 573, 685
 - scalar coupling and, 401
- Amino acid residue
 - chemical shifts in proteins, 783f–785f, 795f–796f
 - random coil, 23, 407f, 408, 782, 794, 804
 - scalar coupling constants for, 800–803
- Amino acid spin system, 787
- ^1H chemical shifts, 784f–785f
- ^1H - ^1H isotropic mixing in, 487–493, 492f, 495f
- HCCH, spin topology, for, 602
- ^{13}C chemical shifts, 795f–796f
- ^{13}C - ^{13}C isotropic mixing in, 611–612, 611f
- triple-resonance, spin topology, for, 615t–617t, 641–642
- Amplifier, rf, 120, 253
- Amplitude modulated signal, 316
- Amplitude, rf, 9–10, 120, 171–173
- Analog signal, 123, 127–129, 156
- Analog-to-digital converter (ADC), 115f, 123, 127–129, 321
- Angle
 - Euler, 104, 107, 108f, 804–805
 - polar, 110, 365, 368, 804–805
- Angular momentum, 2–4 *See also*
 - Magnetic quantum number
 - bulk, 5
 - intrinsic, 29–30
 - nuclear spin, 2–3, 3t
 - orientation of, 3–4
 - quantum mechanical, 29
 - quantum number, 2
 - vectors, 4, 4f
- Angular momentum operator, 46, 80
- commutation relationships for, 46
- eigenfunctions of, 37, 47
- N-spin, 60
- Pauli spin matrix for, 46
- rotation properties of, 51t, 81f
- two-spin, Cartesian basis for, 70
- Anisotropic chemical shift, 112, 668, 751–752, 810–811. *See also*
 - Chemical shift anisotropy
- Anisotropic diffusion. *See* Diffusion
- Anisotropic magnetic susceptibility, 110, 666–667
- Annealing, simulated, 809
- Anticonnected transition, 460
- Antiphase lineshape, 89–90, 413, 413f, 421, 423f, 424–426, 424f, 425f, 470f
- COSY and, 411–412
- representation in spectra, 406
- Antiphase lineshape, double, 443–444, 453, 463, 469–470, 470f, 472t
- Apodization, 138, 143–151. *See also*
 - Apodization function; *individual NMR experiments*
- ^{13}C - ^{13}C scalar coupling, removal of, 655

- Apodization (*Continued*)
 constant time NMR, in, 161, 543–546, 545f
 COSY and, 415–416
 examples of, 146f, 149f, 150f, 417f
 heteronuclear correlation, in, 552
 matched filtering, for, 144, 145, 147–148
 multidimensional NMR, in, 327
 peak integrals influenced by, 148, 151
 resolution enhancement, for, 147–148
 signal-to-noise ratio, and, 158
 truncation artifacts with, 145
- Apodization function
 $\cos^{-1}(\pi J_{CC}t)$, 655
 cosine bell, 145, 146f
 Dolph-Chebyshev, 144
 exponential, 147, 149f, 150f
 Hamming, 145, 146f
 Hanning, 145
 Kaiser, 145, 146f, 149f, 150f
 Lorentzian-to-Gaussian, 147–148, 149f, 150f
 maximum resolution, 148, 149f
 sine bell, phase-shifted, 147, 414
- Apparent scalar coupling. *See* Scalar coupling, apparent
- Aromatic ring, 749
 flip of, 369, 480, 810
 local magnetic fields for, 408
 selective protonation of, 749, 752–753
 scalar coupling constants for, 764
- Aromatic spin system, 480, 485–486
 aliasing and folding for, 444, 483
 chemical shifts of, 782, 783f, 785f, 796f
 decoupling of, 607–608
 fingerprint region for, 418, 419f
 NOE calibration using, 797
- Artifact
 axial peak, 309–310, 323–324
 baseline, 130, 154–158, 156f, 157f, 226, 264f, 265, 298, 307–308, 497
 breakthrough peak, J-filter, 764
 clipped FID, 156–157
 dispersive tails, 411–412, 417f
 double-diagonal, 444–446, 445f
 phase cycle, 310
 pseudo-diagonal, 597
 pulse breakthrough, 161
 repetition rate, 258, 310–311, 444–445
 ridges, 416, 493, 785
 steady-state, 258, 310–311
 t_1 noise, 262, 310, 556
 truncation, 143–148, 146f, 157, 265
 water stripe, 416, 477
- Artifact suppression, 307–310, 313–314
 in heteronuclear correlation, 546–549
 phase cycling, by, 307–310, 313, 437f, 546–549
 pulsed field gradient, by, 313–314, 577f, 584, 598
 spin lock purge pulses, by, 228–230
- Artifact suppression, examples of
 2QF-COSY, 445f, 446f
 4D HMQC-NOESY-HMQC, 598–599
 HSQC, 547–548
- Assignments, resonance. *See* Resonance assignments
- Attenuation, rf power, of, 253
- Audiofrequency filter. *See* Filter, audiofrequency
- Audiofrequency signal, 132
- Autocorrelation function, 138, 348
- Autocorrelation peak. *See* Diagonal peak
- Automated assignment, 791, 794, 809
- Autorelaxation, 339, 340, 357, 359, 362, 375, 381–383, 692. *See also* Cross-relaxation; Relaxation
- AX spin system, 25f, 26, 62–64. *See also* Two-spin system
- Axial peak, 309–310, 323–324

B

- B_0 field gradient. *See* Pulsed field gradient
- B_1 field. *See* rf field
- Back transfer, 507, 527–528
 line, 528
 peak, 527
 plane, 528
- Backbone
 conformation, 806–807
 fingerprint region, 418, 419f
 resonance assignments for, 785–794, 786f, 789f, 790f

- Back-calculation, 810
 - Backprojection (BP), 775
 - Baker-Campbell-Hausdorff (BCH)
 - relationship, 43, 45–46
 - Bandpass filter, 131, 135–136
 - Bandwidth
 - filter, 126
 - noise, 126
 - receiver, 119, 123
 - signal, 124–125
 - Baseline correction, 131, 136, 497
 - Baseline distortion, 130, 154–158, 156f, 157f, 226, 264f, 265, 298, 307–308, 497
 - phase correction and, 156f, 322, 324–326
 - Basic pancreatic trypsin inhibitor (BPTI), 241–242
 - Basis expansion, 32–33, 47, 78–79
 - Basis ket, 38
 - Liouville von Neumann equation and, 41
 - matrix representation and, 47
 - Basis normalization, 30, 79
 - Basis operator space, 78–79
 - Basis operator, 80–83
 - Cartesian, 70, 71t, 80
 - shift, 70, 72, 73t
 - single element, 80, 82
 - single transition, 81, 83f, 100
 - spherical, 70–74, 73t
 - Basis transformation, 47–50, 64
 - Bayesian analysis, 159
 - BCH (Baker-Campbell-Hausdorff relationship), 43, 45–46
 - Biexponential relaxation, 343–345, 681
 - Bilinear operator, 91, 287–288
 - Binomial pulse sequence, 224–227, 227f
 - BioMagResBank, 783, 784f, 794, 795f
 - BIP (Broadband inversion pulse) 179, 197
 - Bloch equations, 7–16
 - chemical exchange and, 392–393
 - derivation of, 7–8, 12–13
 - free-precession, for, 14, 17
 - integration of, 347
 - laboratory reference frame, in, 7–8, 13
 - Larmor frequency in, 8
 - limitations of, 23–24
 - McConnell equations and, 393
 - pulses in, 9, 11
 - relaxation and, 337–338
 - rotating frame, in, 14
 - rotating frame transformation and, 10
 - rotation angle and, 15
 - rotation axis and, 15
 - Bloch vector models. *See* Bloch equations
 - Bloch, Wangsness, and Redfield (BWR), 350. *See also* Relaxation theory
 - Bloch-McConnell equations, 393, 706
 - Bloch-Siegert shift, 9, 170
 - Body diagonal, 527
 - Boltzmann distribution, 5, 12, 333, 335, 350
 - Boltzmann equation, 5, 55, 107
 - Boltzmann probability, 5, 55, 107
 - Bound water, 512–513
 - BP (Backprojection), 775
 - BPTI (Basic pancreatic trypsin inhibitor), 241–242
 - Bra, 37–38, 47, 81
 - Broadband decoupling. *See* Decoupling, phase modulated
 - Broadband inversion pulse (BIP), 179, 197
 - Broadband isotropic mixing. *See* Isotropic mixing
 - Brownian motion, 365, 366. *See also* Diffusion
 - Brownian rotational diffusion, 21
 - Buildup series, 510
 - Bulk angular momentum, 5, 7
 - Bulk magnetic moment, 5, 7, 16
 - Bulk magnetization, 9, 11
 - 180° pulse and, 16
 - equilibrium, 12
 - rf pulse and, 11
 - vector, 9, 12f
 - detection of, 9
 - Burg linear prediction, 161
 - BWR (Bloch, Wangsness, and Redfield), 350. *See also* Relaxation theory
- C**
- $^{13}\text{C}^\alpha$ and ^{13}CO , treated as different nuclear species, 549, 654–655
 - decoupling, 170–172, 549
 - nonresonant phase shift, 170–171, 628

- $^{13}\text{C}^\alpha$ and ^{13}CO (*Continued*)
 rf field strength, adjusting, 169–170, 549
- $^{13}\text{C}^\alpha$ - $^{13}\text{C}^\beta$ decoupling, 549
- ^{13}C chemical shift. *See* Chemical shift, ^{13}C
- ^{13}C isotopic labeling, 533, 727
- ^{13}C - ^{13}C correlation (HCCH), 601–606, 740–743. *See also*
 CC(CO)NH-TOCSY
 HCCH-COSY; HCCH-TOCSY;
 HCC(CO)NH-TOCSY
- $^{13}\text{C}^\beta$ - $^{13}\text{C}^\alpha$ correlation (CBCA), 641–642, 734–735. *See also* CBCANH;
 CBCA(CO)NH; HNCACB;
 HCC(CO)NH-TOCSY;
 CC(CO)NH-TOCSY
- Calbindin $\text{D}_{28\text{k}}$
 CC(CO)NH-TOCSY spectrum, 744f
 description, vi
 ILV NOESY spectrum, 752f
 $^{15}\text{N}/^{15}\text{N}$ separated NOESY spectrum, 747f
 PR HACA(CO)NH spectrum, 774f
 TROSY, multiplet structure in, 572
- Calibration
 NOESY, of, 506–508, 510, 797
 pulse width, 252–257
 temperature, 236–237
- CAMELSPIN (Cross-relaxation
 appropriate for mini-molecules
 emulated by locked spins).
See ROESY
- Capacitance, 119
- Carrier frequency, 9
- Carr-Purcell-Meiboom-Gill (CPMG),
 686–699, 688f, 702, 711–715, 712f,
 720–721, 721f
- Cartesian basis, 70, 71t, 80
- Cartesian operator, 37, 47, 70, 81f, 82,
 90–91
 basis, 70, 71t, 80
 single-element operators and, 82
- Cartesian tensor, 102, 103, 109
- Catenation of 180° pulses, 607, 641
- Causality principle, 142
- CBCACO(N)H, 641
- CBCANH, 617t, 641, 645–650
 product operator analysis for,
 646–648
 pulse sequence for, 648f
 ubiquitin, of, 650f
- CBCA(CO)NH, 616t, 641, 642–645
 pulse sequence for, 643f
 product operator analysis for, 642,
 643–645
 resonance assignment and, 793
 ubiquitin, of, 646
- CC(CO)NH-TOCSY, 741
 calbindin $\text{D}_{28\text{k}}$, of, 744
 pulse sequence for, 742f
- CE-TROSY. *See* Coupling-enhanced
 TROSY
- Characteristic frequency. *See*
 Eigenfrequency
- Chemical exchange, 391–402.
 702–721, *See also* Amide proton
 exchange
 Bloch equations for, 392–393
 chemical shift time scale in, 392,
 399–400, 753–754
 coalescence, in, 399, 754
 CPMG and, 711
 cross-relaxation, similarity to, 397
 fast, 392, 399, 401–402, 754–755
 intermediate, 399, 401, 754–755
 intermolecular, 394, 401, 753–754,
 755f
 intramolecular, 393, 401–402
 isolated spins, for, 392–400
 ligand concentration and, 755f
 lineshape analysis, for, 704–706
 McConnell equations for, 393
 MQ NMR, in, 715–718
 NOESY and, 397, 511
 population transfer and, 395–397,
 397f, 706–707
 relaxation rate constant for, 398–399,
 708, 711, 755f
 ROESY and, 397, 522
 $\text{R}_{1\rho}$ and, 708
 scalar coupled spin systems, 392,
 401–402
 scalar relaxation and, 388
 self-decoupling by, 381
 slow, 392, 399, 401–402, 754–755
 TOCSY, 499
 two-site system, 392, 400f, 755f
 longitudinal magnetization,
 394–397

- rate matrix, 393, 395, 397
- transverse magnetization, 395, 397–399, 708, 711 755f
- ZZ-exchange, 706–707
- Chemical exchange lifetime, 392, 399–400, 753–754
- Chemical kinetics, 392
- Chemical reactions, 391–394. *See also* Chemical exchange
- Chemical shift, 21–22
 - anisotropic, 112, 668, 751–752, 810–811
 - average population weighted, 392, 399
 - ^{13}C resonances, of
 - assignments, use in, 794
 - proteins, in, 795f–796f
 - random coil, in, 794
 - secondary structure dependence, 23, 804
 - conformation-dependent secondary, 23, 782–783, 804
 - degeneracy of, 476, 593
 - 2D NMR, alleviating by, 278
 - 3D heteronuclear-edited NMR, alleviating by, 327–328, 329f, 586f
 - 3D homonuclear NMR, alleviating by, 525, 528
 - 4D heteronuclear-edited NMR, alleviating by, 329–330, 330f, 601, 733
 - triple-resonance NMR, alleviating by, 792
- dispersion, 179, 236, 272, 405, 407f, 408–409, 498, 525, 749, 785, 791
- ethylene glycol, of, 236
- ^1H resonances, of
 - proteins, in, 783f–785f
 - random coil, in, 23, 408, 782
 - secondary structure dependence, 23, 804
- Hamiltonian and, 51, 102, 107
- isotropic, 22
 - dihedral angle restraints from, 804
 - perturbation mapping of, 756–757
- measurement of, 23
- methanol, of, 236
- nuclear shielding and, 22, 102, 107
- perdeuteration and, 732
- referencing of, 23, 262–263
 - indirect, 263t
- tensor, 22, 102, 107
- time scale, 392, 399–400, 753–754
- water, of, 237
- Chemical shift anisotropy (CSA), 22, 383–384, 571. *See also* Anisotropic chemical shift
 - principal axis for, 349–350, 365, 387
 - relaxation, 349–350, 383–384, 502–506, 680
 - relaxation interference, 386–387
 - relaxation rate constants for, 385t
 - spatial functions for, 366t
 - spherical harmonic functions for, 350t
 - tensor operators and, 384t
- Chemical shift asymmetry, 22
- Chemical shift index (CSI), 804
- Chemical shift mapping, 756–757
- Chemical shift time scale, 392, 399–400, 753–754. *See also* Chemical exchange
- CHIRP, 197–198, 210–211
- Circular polarization, 9
- CITY, 493
- Clipped FID, 156–157
- Closure theorem, 38
- ^{13}CO and $^{13}\text{C}^\alpha$, treated as different
 - nuclear species, 549, 654–655
 - decoupling, 170–172, 549
 - nonresonant phase shift, 170–171, 628
 - rf field strength, adjusting, 169–170, 549
- Coalescence, resonance, of, 399, 754. *See also* Chemical exchange
- Cogwheel phase cycling, 307
- CO_H(N)CACB , 641
- Coherence, 70–77. *See also* Double-quantum coherence; Single-quantum coherence; Zero-quantum coherence
 - multiple quantum (MQ), 90–92, 473
 - phase cycling and, 292–295, 298
 - phase shift of
 - frequency discrimination, for, 320
 - rf pulse, by, 298
- pulsed field gradient (PFG), 311
- three-quantum (3Q), 442, 449–551, 481–482
- three-spin single-quantum, 450–451

- Coherence, dephasing of
 pulsed field gradient, by, 217–221, 311
 relaxation, by, 335–336, 346–350
 rf inhomogeneity, by, 227–230, 364, 488
 static magnetic field inhomogeneity, by, 18
- Coherence level. *See* Coherence order
- Coherence level diagram, 293–295
- Coherence order, 293–295
 change in, 294, 298
 heteronuclear, 314, 546–547
 maximum, 293, 307
 relaxation and, 358, 372
 signed, 311, 316
- Coherence order selective (COS), 560.
See also PEP
- Coherence selection, 292–293. *See also* *individual NMR experiments*
 phase cycle, by, 298–307
 pulsed field gradients, by, 311–313, 312f, 438, 574, 576–578
- Coherence transfer, 92–93. *See also* *Magnetization transfer; individual NMR experiments*
 continuous (CW) rf field, by, 286–287
 COSY-type, 281–284, 290–291
 DCOSY, 291–292
 heteronuclear, 290–291
 INEPT, 96–98
 refocused, 98–99, 556f
 in-phase coherence transfer. *See also* *Magnetization transfer*
 COSY-type, 98–99, 283–284, 605–606, 632–633
 DCOSY, 291–292
 TOCSY-type, 284–289, 486–496, 608–612
 isotropic mixing, by, 287–288, 487–493, 492f, 495f, 611f
 NOE, by, 289–290
 pulse-interrupted free precession, by, 280–284
 RDC, by, 291–292
 rf pulse, by, 92, 294
 through-bond, 280
 COSY-type, 281–284, 290–291
 scalar coupling, by, 782
 TOCSY-type, 284–289, 290–291
 through-space, 280
 NOE, by, 289–290
 RDC, by, 291–292
 spin-state selective (S^3CT), 99, 100, 101f
 TOCSY-type, 284–289, 290–291
 triple-resonance, 615t–617t
- Coherence transfer amplitude, 281–282.
See also *individual NMR experiments*
- Coherence transfer pathway, 293–295, 295f. *See also* *individual NMR experiments*
 EXORCYCLE and, 309f
 phase cycle, and, 298–307
 pulsed field gradients and, 311–313, 312f
- Coherent averaging theory, 205
- Coil
 gradient, 217, 219–220, 244
 Helmholtz, 117
 rf, 116f, 117–119, 118f
 shim, 115–117, 115f, 116f, 220, 239–241, 240t, 250–251, 251t
 Z_0 , 117
- Commutation superoperator, 45, 351, 353
- Commutator, 35, 40, 46, 372t
- Commuting operators
 Baker-Campbell-Hausdorff (BCH) relationship for, 43
 exponential matrix of, 43
 simultaneous eigenfunctions, of, 34–35
- Complete relaxation matrix analysis (CORMA), 810
- Complex Fourier transformation, 317, 319, 321. *See also* Fourier transformation
- Composite pulse, 174–179. *See also* *Decoupling*
 180° (π), 174–179, 178f, 203–205
 BIP, 179
 optimal control theory and, 179
- Composite pulse decoupling, 203–209.
See also *Decoupling*
- Computer-aided assignments, 791, 792, 796
- Conformation, protein. *See* Structure determination

- Conformational exchange. *See* Chemical exchange
- Conformation-dependent secondary chemical shift, 23, 804
- Connected transition, 76, 456, 460
- Connectivity, 784–791, 789f, 790f, 792–794
- Constant-time (CT) evolution. *See also individual NMR experiments*
 digital resolution in, 544, 549–550
 evolution period for, 543–544, 549–550
 homonuclear decoupling, 543
 ^{13}C - ^{13}C decoupling in, 578–579
 lineshape in, 543–546, 545f
 ^1H - ^{13}C HSQC
 product operator analysis for, 578–581
 pulse sequence for, 579f
 ubiquitin, of, 581f, 582f
- HSQC
 pulse sequence for, 537
 product operator analysis for, 543–544
 relaxation in, 543–544
 ubiquitin, of, 544f, 545f
 linear prediction and, 161, 162f, 163f
 relaxation in, 543–544
 semi-constant time, 553, 691–692
- Constraint. *See* Restraint
- Continuous Fourier transformation, 136–137, 151. *See also* Fourier transformation
- Continuous wave (CW) rf field, 9, 201–204
- Continuous signal, 124
- Contraction of bra and ket, 37
- Convolution. *See* Apodization
- Convolution difference low pass filter, 232–234, 235f
- Convolution theorem, 137–138, 143–145
- Coordinate system, convention for, 10, 58, 121–122
- CORMA (Complete relaxation matrix analysis), 810
- Correlated Spectroscopy. *See* COSY
- Correlation function. *See* Stochastic function
- Correlation integral, 138
- Correlation theorem, 138
- Correlation time
 effective, 368
 internal, 368
 Hamiltonian, for, 352
 rotational, 18–19, 20f, 21, 336, 352, 363, 366, 729f
- COS (Coherence order selective), 560.
See also PEP
- $\text{Cos}^{-1}(\pi J_{\text{CCt}})$ apodization, 655
- Cosine bell apodization, 145, 146f
- Cosine-modulated signal, 132, 315, 316, 320, 561, 562, 564, 771
- COSY (Correlation Spectroscopy), 409–429. *See also* 2QF-COSY; 3QF-COSY; COSY- β ; DCOSY; DR-COSY; E-COSY; HCCH-COSY; *p*QF-COSY; P-COSY; P.E-COSY; Pre-TOCSY COSY; R. COSY
 apodization for, 147
 cross-peak fine structure in, 412
 experimental protocol for, 412–414
 fingerprint regions in
 aromatic, 418, 419f
 $^1\text{H}^{\alpha}$ - $^1\text{H}^{\beta}$ fingerprint region, 418, 419f
 $^1\text{H}^{\text{N}}$ - $^1\text{H}^{\alpha}$ fingerprint region, 418–419, 419f, 422f
 methyl, 418, 419f, 420f, 421f
 lineshape in, 411–412, 420–422, 423f
 processing of, 415–418
 product operator analysis for, 281–284, 410–412
 pulse sequence for, 410f
 scalar coupling constants, measuring from, 420–426, 423f, 424f, 425f
 self-cancellation in, 284, 414, 420–424, 423f, 424f
 ubiquitin of, 445f, 446f, 449f, 450f, 453f, 454f, 462f, 480f
 variants of, 426–429
- COSY-35, 426–428, 463, 464f
- COSY- β , 426–428, 463
- COSY-type coherence transfer, 281–284, 290–291
- Counter-rotating rf field, 9, 50, 170
- Coupling, dipolar. *See* Dipolar coupling; Residual dipolar coupling
- Coupling, residual dipolar. *See* Residual dipolar coupling

- Coupling, scalar. *See* Scalar coupling
- Coupling-enhanced TROSY (CE-TROSY), 672–673, 673f
- CPMG. *See* Carr-Purcell-Meiboom-Gill
- CRIPT (Cross-relaxation-induced polarization transfer), 726
- Cross-correlation. *See* Relaxation interference
- Cross-peak, 277–278, 277f. *See also individual NMR experiments*
- 3D NMR and, 529f, 586f
- amino acid spin system and, 782
- fine structure
- COSY, 412, 414
- E.COSY, 456f, 459f
- pQ NMR, 469–473
- pQF-COSY, 452, 453–455, 456f
- forbidden, 453, 483
- homonuclear 3D, in, 527–528
- Cross-polarization, 280, 286, 290. *See also* Isotropic mixing; TOCSY
- Cross-rate constant, 342
- Cross-relaxation, 340, 346, 357, 362–363, 372–375, 680–685. *See also* NOE; NOESY; ROE; ROESY
- dipolar, 20f, 359–360, 365–366, 370–383, 377t. 346, 390, 797
- forbidden cross-peaks and, 360, 453, 483
- laboratory frame, 357, 359, 362, 374–375, 390
- NOE and ROE, compensating for, 391, 491, 493
- rotating frame, 381–383, 390
- secular approximation and, 358, 360–363, 372, 376, 380, 683
- through-space correlation by, 289–290
- Cross-relaxation appropriate for mini-molecules emulated by locked spins (CAMELSPIN). *See* ROESY
- Cross-relaxation-induced polarization transfer (CRIPT), 726
- Cross-saturation, 757–759, 758f
- Cross-spectral density function, 359, 369–370
- Cryogenic probe, 119–120, 222, 252
- CSA. *See* Chemical shift anisotropy
- CSA relaxation. *See* Chemical shift anisotropy
- CSA/dipolar relaxation interference, 359–360, 373, 385–387, 570–572, 672, 686, 692, 695f, 703–704, 713
- CSI (Chemical shift index) 804
- Curvature, baseline, 154–158, 156f, 157f
- CW (continuous wave) rf field, 9, 201–204
- Cyanometmyoglobin, 667
- Cyclically ordered phase sequence (CYCLOPS), 266, 307–308
- Cycling sidebands, 203, 211–217, 214f, 215f
- CYCLOPS (Cyclically ordered phase sequence), 266, 307–308

D

- D₂O, 21, 222–223, 224f, 259, 260f, 406, 420, 444, 461, 474, 500, 573, 588, 727, 753, 759, 806
- Damped oscillator, 139. *See also* FID; Interferogram
- Data processing, 129, 136–165, 263–267
- HSVD and, 161
- LPSVD and, 161
- NMR spectrometer, in, 136
- DCOSY (Dipolar correlation spectroscopy), 292
- Decay constant, 139, 399, 400, 692
- Decibel (dB), 253
- Decimation, 129
- Deconvolution of N/P signals, 576
- Deconvolution of orthogonal signals, 562, 564, 567, 576
- Decoupled HSQC. *See* HSQC, decoupled
- Decoupled inversion recovery, 345, 346f
- Decoupling, 96, 98, 181, 201–217. *See also* Isotropic mixing; *individual decoupling sequences*
- 180° pulses, by, 96, 201, 203, 204
- HSQC, in, 541, 547–548
- selective, 170–172, 549, 579
- adiabatic, 209–212
- antiphase magnetization, effect on, 98–99, 556
- asynchronous, 203
- composite pulse, 203–209
- constant-time evolution, in, 543, 578–581

- cycling sidebands, 212–216
- ^1H , 552–553, 555–557, 570f, 626–627
- HSQC, in, 555, 697
- omitting, effects of, 596, 618
- recommendations for, 216–217
- relaxation and, 364
- scalar coupling and, 96, 98
- scaling factor, 204
- selective, 271
- self-, 381, 660, 799
- semiselective, 549
- supercycle, 205
- synchronous, 555, 627
- theory of, 201–204
- Decoupling in the presence of scalar interaction. *See* DIPSI
- Degenerate NOE, 515, 516f, 593, 598
- Degeneracy, resonance. *See* Chemical shift
- Degenerate transition, 357, 358f, 361
- Denaturation, 407f, 408, 409
- Density matrix, 37, 41. *See also* Density operator
 - BCH and, 43, 45
 - closure theorem and, 38
 - density operator and, 41
 - diagonal elements, 55, 74, 333
 - Dirac notation and, 37–38
 - off-diagonal elements, 74
 - rotating frame transformation and, 43
 - similarity transformation and, 44
 - trace of, 39
- Density matrix formalism, 29, 78, 401
- Density operator, 41, 68, 361
 - BCH and, 43, 45
 - diagonal elements of, 333
 - equilibrium, 55–56, 333
 - high temperature approximation for, 55
 - semi-classical relaxation, in, 352
 - two-spin system, for, 68
 - expansion of, 77–80, 294
 - interaction frame, in, 351
 - Liouville von Neumann equation and, 42
 - off-diagonal elements of, 333, 336
 - relaxation of, 351–359
 - rotating frame, in, 43
 - time-evolution of, 41–43, 280
 - transformation of, 79
- Dephasing, coherence, of. *See* Coherence, dephasing of
- DEPT (Distortionless Enhancement by Polarization Transfer), 290
- Derivative theorem, 137
- Detected nucleus, 535
- Detection, indirect, 535
- Detection operator, 55–56, 89, 308
- Detection, phase sensitive. *See* Frequency discrimination; Quadrature detection
- Detection, proton, 535
- Detection, quadrature. *See* Frequency discrimination; Quadrature detection
- Detection, single-channel, 132
- Detector, phase sensitive, 120–123, 132–136
- Deuteration, 533, 726–728, 730f, 733, 748, 759
 - CT-HNCA and, 735–737
 - experiments in, 733–744
 - HNCACB and, 737–739
 - HNCO and, 739
 - HN(COCA)NH and, 739
 - NOE restraints from, 745–749
 - relaxation and, 730f
 - side chains and, 740–743
 - static magnetic field and, 739
- Deuteration, complete. *See* Perdeuteration
- Deuteration, fractional, 533, 727–733, 730f, 737, 740, 745
- Deuterium, 117, 222, 727, 748
 - decoupling, 697, 734, 735, 749
 - isotope shifts and, 732f, 733
 - lock circuit, 117
 - relaxation rate constants and, 693–694
 - shimming and, 241, 252
- Deuteron, 2, 222, 500, 697, 728, 732, 734. *See also* Deuterium
- DG (Distance geometry), 808
- DHPC. *See* Dihexanoyl phosphatidylcholine
- Diagonal
 - anti-, 520
 - double, 445f, 446f
 - pseudo-, 471f, 475f, 597

- Diagonal peak, 277, 290, 452–453, 475, 593, 597
- Diagonalization, 63, 90, 341
exponential matrix, of, 341
Hamiltonian, of, 50, 63
- Diamagnetic molecule, 102, 111, 370
- Diamagnetic protein, 667, 668
- Differential equation
coupled, 357
evolution in time, 41
first order, 339, 341
harmonic, 84
integration of, 352
- Diffusion. *See also* Spin diffusion
Brownian rotational, 21, 335, 365–367
rotational
anisotropic, 19, 367, 370
Brownian motion, in, 21, 335, 365–368
correlation time, for, 19–21, 352, 366, 370
isotropic, 19–21, 365–367
relaxation, by, 352, 366
- Diffusion, spin. *See* Spin diffusion
- Digital filter, in NMR spectrometer, 126–131
- Digital phase shifting, 438
- Digital Quadrature Detection (DQD), 136
- Digital resolution. *See* Resolution, digital
- Digital resolution enhancement, 149f, 150f
- Digital signal processing, 124, 129–130, 135, 142, 223
- Digitizer, 123, 128, 133f
- Digitizer phase, 298, 308
- Dihedral angle, 455, 656
restraint, 798–804
Karplus, curve, dependence on, 388, 798–803
time-dependent, 388, 799–800
- Dihedral angle restraint
from isotropic chemical shift, 804
from scalar coupling constant, 798–803
- Dihexanoyl phosphatidylcholine (DHPC), RDC and, 667
- 2,2-dimethyl-2-silapentane-5-sulfonic acid (DSS), 262–263, 263t
- Dimyristoyl phosphatidylcholine (DMPC), RDC and, 667
- Dioxane, 262
- Dipolar correlation spectroscopy (DCOSY), 292
- Dipolar coupling, 24
coherence transfer under, 291
Hamiltonian and, 80, 110–111
heteronuclear, 290–291
homonuclear, 289
like spins, for, 381–383
magnetization transfer by, 289, 346, 502–510
measurement of, 665
principal orientation of, 365
spatial variables for, 366t
spherical harmonic functions for, 350t
tensor operators for, 372t, 382
- Dipolar coupling constants, restraints from, 804–805
- Dipolar field, 371
- Dipolar-CSA relaxation interference, 359–360, 373, 385–387, 570–572, 672, 686, 692, 695f, 703–704, 713
- Dipolar relaxation, 20f, 359–360, 365–366, 370–383, 377t, 386, 502, 571–572, 680, 685 *See also* Cross-relaxation; NOE; NOESY; ROE; ROESY
- Dipole-dipole (DD) relaxation. *See* Dipolar relaxation
- DIPSI (Decoupling in the presence of scalar interaction), 364, 490–493
HCCH-TOCSY and, 610
HSQC and, 555
pulse sequences and, 490
- DIPSI-2, 364, 490–493
- DIPSI-2 relaxation-compensated (DIPSI-2rc) sequence, 486, 493, 496, 515, 590
- DIPSI-2rc. *See* DIPSI-2 relaxation-compensated sequence
- DIPSI-3, 611
- Dirac notation, 37
basis operators and, 80
closure theorem and, 38
density matrix and, 37–38
Liouville von Neumann equation and, 41

- Schrödinger equation and, 41
 - Direct peak, 469
 - 2Q and, 471, 475f, 479f
 - 3Q and, 484f, 486, 487f
 - Direct product, 59–61
 - Discrete Fourier transformation, 140–143, 154, 158. *See also* Fourier transformation
 - Discrete signal, 124
 - Dispersion, chemical shift, 179, 236, 272, 405, 407f, 408–409, 498, 525, 749, 785, 791
 - Dispersive lineshape, 18, 139–140, 140f, 317
 - Dispersive antiphase lineshape, 425f, 469, 470f, 471, 474, 476–477
 - Dispersive tails, 235f, 319, 412, 417f, 431, 434f, 442, 444, 471, 475, 490, 505
 - Displacement vector, 459
 - Distance constraint, 507–508, 510, 522. *See also* Distance restraint
 - Distance geometry (DG), 808
 - Distance, lower bound, 798
 - Distance, upper bound, 510, 522, 797, 798
 - Distance restraint, 499, 502, 510, 527, 745, 749, 751–752, 756, 760, 762, 796–798, 805, 807–808. *See also* Distance constraint
 - Distortionless Enhancement by Polarization Transfer (DEPT), 290
 - DMPC. *See* Dimyristoyl phosphatidylcholine
 - Dolph-Chebyshev window, 144–145
 - Double difference isotope filter, 547
 - Double quantum NMR. *See* 2Q NMR
 - Double-half-filter. *See* Isotope filter
 - Double-quantum (DQ, 2Q) coherence, 74, 75, 82, 83f, 101f, 293, 295f, 301, 312, 441, 443, 480, 697. *See also* 2Q NMR; Coherence
 - bilinear product operators and, 90–91
 - chemical exchange and, 715
 - chemical shifts and, 91
 - HMQC and, 540, 718
 - NOE and, 505
 - selection of, 301f, 302t
 - Double-quantum filtered COSY. *See* 2QF-COSY
 - Double-quantum (DQ) splitting, 468–469
 - Double-quantum (DQ) transition, 76, 82–83, 346
 - Double-relayed COSY. *See* DR.COSY
 - Doubly absorptive lineshape, 319
 - Doubly antiphase absorptive lineshape, 415, 470, 483
 - Doubly antiphase dispersive lineshape, 469–471, 477
 - Doubly dispersive lineshape, 319
 - Doubly rotating tilted frame, 363–364, 381–383
 - DQ. *See* Double-quantum (DQ) coherence
 - DQD (Digital Quadrature Detection), 136
 - DR.COSY (Double-relayed COSY), 433–436, 785
 - absorptive antiphase lineshape in, 434
 - pulse sequence for, 435f
 - ubiquitin, of, 436f
 - DSS (2,2-dimethyl-2-silapentane-5-sulfonic acid), 262, 263
 - Dynamic frequency shift, 356–357
 - Dynamic range, 123, 128, 135, 156, 203, 221, 223, 265
 - Dynamical simulated annealing (SA), 809
 - Dynamics, internal, 334, 365–370, 409, 499
 - aromatic ring flips, 369, 810
 - chemical shifts, effect on. *See* Chemical exchange
 - methyl rotation, 369, 406, 810
 - NOE, effect on, 369
 - scalar coupling, effect on, 799, 802
- ## E
- E.COSY (Exclusive correlation spectroscopy), 455–463
 - displacement vector, 459
 - experimental protocol for, 460–461
 - heteronuclear, 656–657, 660, 670
 - lineshape asymmetry, 660
 - phase cycles for, 461t
 - processing of, 461–462

- E.COSY (*Continued*)
 product operator analysis for, 457–458
 self-decoupling in, 660
 ubiquitin, of, 462f, 464f, 661f
 variants of, 463
 weight factors for, 457
- Eigenbra, 46–47
- Eigenfrequency, 353, 360
- Eigenfunction, 31
 Hamiltonian, of, 50
 Hamiltonian commutation superoperator, of, 353
- Heisenberg uncertainty relationship and, 35
- Hermetian operator, of, 32
- simultaneous, 34–35
- stationary state, 34
- strong coupling Hamiltonian, for, 62–63
- Eigenket, 38, 46–47
- Eigenoperator, 353–354, 361
- Eigenstate, 63, 74, 81–82. *See also* Stationary state
 coherence order and, 293–294
- Eigenvalue, 31–34, 341, 362, 395, 398
- Eigenvalue equation, 31–34
- Electric field gradient tensor, 365, 366t, 384
- Electric quadrupole moment.
See Quadrupole moment
- Energy, 4, 31
- Energy level, 4–7, 24–26, 62–64, 82–83, 335, 338
- Enhancement, NOE, 344–345, 389–391
- Ensemble of 3D structures, 808–813, 811
- Envelope function, 144
- Equilibrium
 chemical, 393
 dissociation constant, 757
 orientational, 106–107, 368
 thermal
 density operator, 55–56, 75, 294
 lattice, of, 335–336
 magnetization, 5, 17f
 phase cycling and, 294, 306, 310–311
 relaxation to, 257–258, 333, 336, 350, 352
- Ernst angle, 257–258, 258f
- Ethylene glycol, temperature calibration using, 236
- Euler angles, 104, 107, 108f, 804–805
- Evolution. *See* Liouville von Neumann equation; Product operator formalism
- Evolution period, indirect. *See* Indirect evolution period
- Exchangeable proton. *See* Amide proton exchange
- Excitation null, 169–170
- Excitation sculpting, 230, 233f
 in 3D NOESY-TOCSY spectra, 526f
 in NOESY, 504f
 in TOCSY, 489f
- Exclusive correlation spectroscopy.
See E.COSY
- Exclusive COSY. *See* E.COSY
- EXORCYCLE, 266, 309, 313
- Expectation value, 33–35, 40–41, 48, 79
 of magnetic moment, 35–37
- Exponential apodization, 147, 149f, 150f
- Exponential of matrix, 341
- Exponential operator, 42–43, 45–46
- Extreme narrowing, 367, 384, 391

F

- F_1 , 276
- F_2 , 274
- Faraday's law, 9
- Fast exchange, 392, 399, 401–402, 754–755. *See also* Chemical exchange
- Fast Fourier Transformation (FFT), 142. *See also* Fourier transformation
- FFT (Fast Fourier Transform), 142.
See also Fourier transformation
- FID (Free induction decay), 17, 19f, 134f
 clipping of, 156–157
 distortion by filter response, 127, 157
 interferogram and, 274
 quantization noise in, 127–129
 shimming, used for, 241
- Field gradients. *See* Pulsed field gradients

- Filamentous phage, RDC and, 668, 752
- Filter. *See also* Isotope filter
- analog, 126–127
 - audiofrequency, 115f, 123, 132–133, 133f, 157
 - baseline distortions, 156–158, 493–494
 - phase delay, 127
 - transient response, 127, 131, 157, 266
 - bandpass, 135–136
 - digital, 129–131, 232–234, 235f
 - first-order J , 767
 - half, 763–769, 766f, 766t
 - low-pass, 133, 235f
 - second-order J , 764, 768
- Filter diagonalization, 770
- Filter function. *See* Apodization function
- Filter, isotope. *See* Isotope filter
- Filtering. *See* Apodization
- Fine structure. *See* Cross-peak, fine structure
- Fingerprint region, 418–421
- aromatic, 418, 419f
 - $^1\text{H}^\alpha$ - $^1\text{H}^\beta$ fingerprint region, 418, 419f, 453–455, 454f, 455f, 478, 479f
 - $^1\text{H}^N$ - $^1\text{H}^Z$ fingerprint region, 418–419, 419f, 422f
 - methyl, 418, 419f, 420f, 421f, 453f
- Finite impulse response (FIR), 129
- FIR (Finite impulse response), 129
- First-order phase correction, 153f
- Flip angle, pulse, 11, 52
- Flip-back, 231–232, 574–578, 575f
- saturation transfer and, 690–691
- Flip-flip transition, 75, 76f, 83f, 339, 346
- Flip-flop spectroscopy. *See* FLOPSY
- Flip-flop transition, 75, 76f, 83f, 339, 346
- FLOPSY (Flip-flop spectroscopy), 490–491, 492f
- HCCH-TOCSY and, 610–611
- FM (Frequency modulation frame), 191–192, 196
- Folding. *See* Aliasing and folding
- Forbidden cross-peak, 360, 453, 483
- Force field, molecular dynamics, 808–809
- Four-dimensional NMR. *See* 4D NMR
- Fourier analysis, 175, 180, 335
- Fourier transform algorithms
- complex, 317, 319, 321
 - continuous, 136–137, 151
 - discrete, 141–142, 151, 154, 158
 - fast (FFT), 142, 770
 - inverse, 137, 600t, 661f
 - linearity of, 137
 - periodicity of, 125
 - real, 141, 317, 318f, 319, 321–322
 - scaling first point in, 156
- Fourier transform theorems, 137–138
- Fourier transformation, 131
- alternatives to, 159–165
 - apodization in, 143
 - baseline roll and, 157
 - causality in, 142–143
 - damped oscillator, of, 139, 142, 316–317
 - Dolph-Chebyshev window and, 144
 - FID, of, 17–18, 19f
 - frequency discrimination and, 318f
 - Lorentzian lineshape and, 140f, 317
 - nonuniform sampling and, 770–771
 - sinc-wiggles with, 157
 - truncation artifacts and, 143–148, 146f, 157, 265
 - zero-filling in, 142
- Fractional deuteration, 533, 727–733, 730f, 737, 740, 745
- Free induction decay. *See* FID
- Free precession, 14, 17
- coherence order, conservation of, 294
 - evolution in the product operator formalism, 84–85
 - Hamiltonian, 61–62, 64, 538, 602
 - pulse-interrupted, 280–284
- Frequency
- audio, 132, 135
 - carrier, 9
 - intermediate, 135
 - Larmor, 8
 - Nyquist, 124–126, 126f
 - transmitter, 9
- Frequency, characteristic. *See* Eigenfrequency

Frequency discrimination, 132–135, 133f, 134f, 315–326, 318f. *See also* Quadrature detection
 hypercomplex (States), 315, 320–321, 323t
 N/P selection for, 316–317, 322, 576
 PEP (preservation of equivalent pathways), 560–566, 574–578
 phase cycling, by, 320–322
 p Q NMR, in, 473
 pulsed field gradient, by, 322–323, 574–578
 Redfield's method for, 321
 time-proportional-phase-incrementation (TPPI), 315, 321–322, 323t
 TPPI-States, 322, 323t, 324–326, 325t, 655
 Frequency labeling, 279–280
 Frequency modulation (FM) frame, 191–192, 196
 Frequency, resonance, 8, 11, 125 132, 237, 324
 Frequency shift
 dynamic, 356–357
 nonresonant, 9, 170–172
 off-resonance, 9, 170–172
 Frequency-shifted pulse, 181–182, 184–189
 Frequency-shifting theorem, 137–138
 Full-width-at-half-height (FWHH), 18, 139–140, 147–148, 408, 703
 FWHH. *See* Full-width-at-half-height

G

GARP (Globally Optimized Alternating-phase Rectangular Pulse), 206, 208f, 214–215, 214f, 217, 690
 Gaussian lineshape, 148, 149f, 150f
 Gaussian selective pulse, 182f, 183f, 184f
 Geminal proton
 NOE calibration, for, 797
 spin-diffusion in, 522
 GFT-NMR (G-matrix Fourier Transform NMR spectroscopy) 771–772
 Global fold, 749, 751, 786f, 807, 807f
 Globally Optimized Alternating-phase Rectangular Pulse. *See* GARP

G-matrix Fourier transform (GFT) NMR spectroscopy, 771–772
 Gradient echo, 218, 230, 311–312, 768
 Gradient enhancement, heteronuclear correlation and, 574–578
 Gradient, pulsed field. *See* Pulsed field gradient
 Gradient shimming, 242, 244–252
 3D profile edge shimming 252
 pulse sequence for, 248f
 shim map for, 251f
 Z1 profile shimming, 252
 Gradient-enhanced (PFG) HNCA, 620f, 627–628
 Gradient-enhanced (PFG) HSQC, 574, 576–578, 577f
 Gradient-enhanced (PFG) TROSY, 574, 576–578, 577f
 Gradient-enhanced (PFG) TROSY-HNCA, 620f, 628
 Gram-Schmidt process, 32
 Group theory, 62, 361, 373

H

^1H 3D NMR, 525–529
 experimental protocol for, 526–527
 processing of, 527
 pulse sequence for, 526
 variants of, 528–529
 ^1H chemical shifts. *See* Chemical shift, ^1H
 ^1H decoupling, 552–553, 555–557, 570f, 626–627
 ^1H detection, 535
 ^1H linewidth, 406, 408, 407f
 ^1H NMR spectrum. *See* 1D ^1H NMR
 ^1H relaxation rate constants, 20f, 729f, 730f
 ^1H resonance assignments, 782–791
 $^1\text{H}^\alpha$ - $^1\text{H}^\beta$ fingerprint region, 418, 419f, 453–455, 454f, 455f, 478, 479f
 $^1\text{H}^\text{N}$ - $^1\text{H}^\alpha$ fingerprint region, 418–419, 419f, 422f
 ^2H . *See* Deuterium
 HACANH, 774f, 775
 HACA(CO)NH, 774f, 775
 Hahn echo, 264f, 265–266, 267f, 704f
 2D NMR, in, 488f, 503f
 baseline distortions, 264f, 265–266

- chemical exchange and, 718, 719f
- jump-return and, 261f
- phase, 265–266, 264f
- pulse sequence for, 228f, 229f, 265–266, 704f, 719f
- spin-spin relaxation, for, 259, 261f
- solvent suppression, 228f, 229f, 232, 263–267
- ubiquitin, of, 267f, 407f, 783f
- Half-filter. *See* Isotope filter
- Half-Gaussian selective pulse, 182f, 183f
- Hamiltonian, 30, 34, 102–104, 106, 351
 - averaging of, 102–112
 - chemical shift, 102, 107, 349
 - commutation superoperator, 351, 353
 - diagonal, 49–50
 - diagonalization, of, 49–50, 62–63
 - dipolar coupling
 - principal orientation of, 365
 - spatial variables for, 366t
 - spherical harmonic functions for, 350t
 - tensor operators for, 372t, 382
 - residual, 80, 110–111, 291
- effective, 44–45, 50–51
- eigenbase and, 43, 55. *See also* Eigenfunction
- eigenfunction of, 34, 49–50
- eigenvalue of, 34, 49–50
- free precession, 61–62, 84, 602
- interaction frame, in, 351–352, 363
- isotropic chemical shift, 51, 84, 107
- in isotropic phase, 107
- Liouville von Neumann equation and, 42
- magnetic field, 50
- pulse, 85–86
- quadrupolar, 350t, 366t, 385t, 680
- relaxation and, 351
- residual dipole coupling, 80, 110–111, 291
- rf, 50, 85–86
- rotating frame, in, 43–45, 363
- scalar coupling and, 61–67, 107, 287
 - strong, 26, 61–64, 90, 284
 - weak, 64, 84
- similarity transformation of, 44–45
- stationary states of, 34, 49–50
- stochastic, 335, 351, 353
- strong coupling, 61–62
- time dependent, 351
- time independent, 31, 34, 42, 351
- transformation under rotation of, 104, 108
- truncation of, 104, 106, 108
- weak coupling, 64, 84
- Zeeman, 50, 57, 363
- Hamiltonian superoperator, 351, 353
- Hamming apodization, 145, 146f, 148
- Hankel singular value decomposition (HSVD), 161, 162f, 163f, 498
- Hanning apodization, 145
- Harmonic signal, 132
- Hartmann-Hahn condition, 286–289.
See also Isotropic mixing; TOCSY
- isotropic mixing sequences and, 288
- HBHACBCA(CO)NH, 739
- HBHACBCANH, 739
- HBHA(CBCACO)NH, 645
- HBHA(CBCA)NH, 649
- HBLV (Hybrid Backprojection Lower Valve), 775
- HCA II (Human carbonic anhydrase II), 731–732
- HCACO, 637
- H(CA)NH, 615t, 632–636
 - product operator analysis for, 633–636
 - pulse sequence for, 633f
 - ubiquitin, of, 636f
- (H)CC(CO)NH-TOCSY, 741, 743
- HCCH-COSY, 601–608
 - constant-time
 - pulse sequence for, 603f
 - product operator analysis for, 607–608
 - ubiquitin, of, 609f
 - product operator analysis for, 604–607
 - pulse sequence for, 603f
- HCCH-TOCSY, 601–603, 608–612
- Isotropic mixing in, 611f
- product operator analysis for, 608–610
- pulse sequence for, 610f
- ubiquitin, of, 612f
- HCC(CO)NH-TOCSY, 743
- Heat bath. *See* Lattice
- Heisenberg uncertainty principle, 35, 336

- Helmholtz coil, 117
- Hermitian operator, 31–32, 39, 41
- Heteronuclear coherence order, 314, 546–547
- Heteronuclear coherence transfer, 290–291
- Heteronuclear correlation NMR, 533–581. *See also individual NMR experiments*
- aliasing and folding, in, 546, 549–552, 550f
- artifact suppression in, 546–549
- ^{13}C - ^{13}C scalar coupling in, 548–549
- isotope filter, in, 546–547
- gradient enhancement and, 573–578
- initial sampling delay in, 325–326
- phase cycling and, 546–549
- processing of, 552
- quadrature detection and, 546
- sensitivity of, 534–535
- solvent suppression in, 573–574
- Heteronuclear E.COSY, 656–660, 661f, 670
- Heteronuclear Multiple-Quantum Coherence. *See* HMQC
- Heteronuclear NMR experiments, 533
- Heteronuclear scalar coupling constants, 617t
- Heteronuclear single-quantum coherence. *See* HSQC
- Heteronuclear-edited NMR, 581–601, 745–753. *See also individual NMR experiments*
- Heteronuclear-edited NOESY. *See individual NMR experiments*
- Heteronuclear-edited TOCSY. *See individual NMR experiments*
- High resolution 3D structure, 806–813
- High temperature limit, 5, 56, 356, 365
- Hilbert space, 32, 55
- Hilbert transform, 143, 357
- HMQC (Heteronuclear Multiple-Quantum Coherence), 535–540. *See also* HSQC, TROSY
- chemical exchange and, 718
- EXORCYCLE and, 547
- isotope filter in, 546–547
- jump-return in, 226
- lineshape in, 536, 538, 546, 549
- phase cycle for, 546–547
- processing for, 552
- product operator analysis for, 536, 538–539
- pulse sequence for, 537f
- relaxation in, 539–540, 545–546, 545f
- ubiquitin, of, 544f, 545f
- HMQC-NOESY-HMQC, 593–601, 762. *See also* HMQC-NOESY-HSQC; HSQC-NOESY-HSQC
- $^{13}\text{C}/^{13}\text{C}$, 597–599, 598f, 600t
- $^{13}\text{C}/^{15}\text{N}$, 595–597, 596f
- $^{15}\text{N}/^{15}\text{N}$, 594–595, 594f
- processing, 599–600, 600t
- pulse sequence for, 594f, 596f, 598f
- HMQC-NOESY-HSQC, 597
- HNCA, 614, 615t, 618–628, 734–737
- constant-time, 625–626
- decoupled, 626–627
- deuteration and, 735–737
- DQ, 740
- gradient-enhanced, 627–628
- HN(CO)CA and, 629, 740
- intraresidue and interresidue connectivities, distinguishing, 740
- ^{13}C - ^{13}C scalar coupling in, 622–623
- pulse sequence for, 619f–620f, 736f
- product operator analysis for, 618, 621–627
- relaxation in, 623, 625–626
- sensitivity of, 625–626
- TROSY, 628
- ubiquitin, of, 624f
- HNCACB, 641–642, 650–654. *See also* CBCANH, HN(CA)CB
- pulse sequence for, 651f, 738f
- product operator analysis for, 651–653
- resonance assignment and, 793–794
- ubiquitin, of, 654f, 774f
- HN(CA)CB, 737–739. *See also* CBCANH, HNCACB
- deuteration and, 737–739
- pulse sequence for, 738f
- HN(CACB)CG, 740
- HN(CA)CO, 616t, 639–641
- pulse sequence for, 640f
- sensitivity of, 641

- HNCA-J, 656–660
 $^3J_{H^N H^\alpha}$ measured by, 660, 665, 667t
 pulse sequence for, 657f
 product operator analysis for, 657–660
 relaxation in, 660
 ubiquitin, of, 661f, 667t
- HNCO, 637–638
 product analysis for, 637–638
 pulse sequence for, 638f
- HN(CO)CA, 629–632
 intraresidue and interresidue connectivities, distinguishing, 740
 product operator analysis for, 629–631
 pulse sequence for, 630f
 of ubiquitin, 632f
- HN(CO)CACB, 641. *See also* HN(COCA)CB
 resonance assignment and, 793–794
- HN(COCA)CB, 734–735, 738–739
- HN(COCA)NH, 739
- HN(CX)_nCY, 740–741
- HNHA, 660, 662–665
 $^3J_{H^N H^\alpha}$ measured by, 664–665, 667t
 product operator analysis for, 662–664
 pulse sequence for, 662f
 relaxation in, 664, 665f
 ubiquitin of, 666f, 667t
- HOHAHA. *See* Homonuclear Hartmann-Hahn
- Homogeneity, magnetic field, 18, 115–116, 223, 238–239, 241, 245–247
- Homonuclear 3D NMR. *See* ^1H 3D NMR
- Homonuclear chemical shifts. *See* Chemical shifts, ^1H
- Homonuclear Hartmann-Hahn (HOHAHA), 280–281, 286, 288–289, 486. *See also* isotropic mixing; TOCSY
- Homonuclear scalar coupling
 constants, 401, 504, 535, 538, 539, 541, 543, 583, 670, 703, 714, 726, 734, 798–802, 800f, 800t, 801t, 802f, 802t, 803f, 803f, 806–807
- Homospoil pulse, 220, 574
- HSQC (Heteronuclear single-quantum correlation), 535. *See also* HMQC; TROSY
 ^{13}C scalar coupling and multiplet structure, 548–549
 chemical exchange and, 718
 chemical shift mapping and, 756
 constant-time, 543–544
 pulse sequence for, 537
 product operator analysis for, 543–544
 relaxation in, 543–544
 ubiquitin, of, 544f, 545f
- decoupled, 553–560
 energy level diagram for, 559f
 pulse sequence for, 554f
 product operator analysis for, 553, 555–560
 relaxation in, 557–558
 ubiquitin, of, 570f
- fast, 573–574, 575f
- folding and aliasing in, 449–452, 550f, 551f
- ^1H - ^{13}C
 product operator analysis for, 578–581
 pulse sequence for, 579f
 ubiquitin, of, 581f, 582f
- interferogram for, 545f
- IPAP, 670–672, 671f
- phase cycling and artifact suppression, 546–548
- processing, 552
- product operator analysis for, 540–542
 pulse sequence for, 537f, 554f
- RDC measured from, 669–670, 669f
- relaxation in, 338, 542, 543–544, 557–558, 562–563
- sensitivity-enhanced (PEP), 560–566
 energy level diagram for, 559f
 pulse sequence for, 554f
 product operator analysis for, 560–566
 relaxation in, 562–563
 ubiquitin, of, 570f
- ubiquitin of, 162f, 163f, 165f, 544f, 545f, 551f, 581f

- HSQC-NOESY, 591–593, 762
 pulse sequence for, 592f
 HSQC-NOESY-HSQC, 745–753. *See also* HMQC-NOESY-HMQC;
 HMQC-NOESY-HSQC
 $^{15}\text{N}/^{15}\text{N}$ -separated, 745–747
 pulse sequence for, 746f
 calbindin D_{28k}, of, 747f, 752f
 $^{13}\text{C}/^{15}\text{N}$ -separated, 747–749
 pulse sequence for, 750f
 calbindin D_{28k}, of, 752f
 $^{13}\text{C}/^{13}\text{C}$ -separated, 747–749
 calbindin D_{28k}, of, 752f
 HSQC-TOCSY, 591–593
 HSSVD (Hankel singular value
 decomposition) linear prediction,
 161, 162f, 163f, 498
 Human carbonic anhydrase II (HCA II),
 731–732
 Hybrid Backprojection Lower Valve
 (HBLV), 775
 Hydration layer, 21
 Hydrodynamic radius, 21
 Hydrogen bond restraints
 from amide proton-solvent exchange,
 805–806
 from trans-hydrogen bond scalar
 coupling constants, 806
 Hydrogen exchange. *See* Amide proton
 exchange
 Hydroxyl proton, 512–513
 Hypercomplex (States) frequency-
 discrimination, 315, 320, 322–325,
 511, 550, 564, 576
- I**
- Identity operator, 38, 55–56
 IF (Intermediate frequency), 135
 Impedance, 118–119, 237
 Impurity, detecting, 409
 Incrementable delay, 273, 279, 326–327.
See also Indirect evolution period
 Indirect detection, 535
 Indirect evolution period, 274, 275f,
 278–280. *See also* Acquisition
 period
 constant time, 543–541, 578–581
 initial delay, for, 172, 325–326, 546
 semi-constant time, 553, 691–692
 Indirect evolution time, 273, 279,
 326–327. *See also* Indirect
 evolution period
 Indirect magnetization transfer. *See*
 Spin diffusion
 Inductance, 119
 INEPT (Insensitive Nuclei Enhanced
 by Polarization Transfer), 96–99.
See also PEP; TROSY
 homonuclear scalar coupling in,
 541–542
 HSQC and, 540–542, 543, 553
 optimal transfer, 338
 refocused, 98–99, 553, 554f, 555–560,
 556f, 633
 relaxation in, 338
 solvent suppression in, 573–574
 Inhomogeneous broadening, 18,
 534–544, 545f
 Initial rate approximation, 343–345, 507,
 681, 797
 Initial rate regime, 343–345, 507, 681,
 797
 In-phase coherence transfer. *See also*
 Magnetization transfer
 COSY-type, 98–99, 283–284,
 605–606, 632–633
 DCOSY, 291–292
 TOCSY-type, 284–289, 486–496,
 608–612
 In-phase lineshape, 88–90
 I_nS spin system, 553, 555–557, 556f,
 598
 Insensitive Nuclei Enhanced by
 Polarization Transfer. *See* INEPT
 Integrated intensity, 148, 151, 505–506
 Interaction frame, 351–352
 Interference, between relaxation
 interactions. *See* Relaxation
 interference
 Interference, destructive, 284, 453, 508,
 572
 Interference effects. *See* Relaxation
 interference
 Interferogram, 124, 144–145, 274–276,
 275f. *See also* FID
 heteronuclear correlation and, 545f
 Intermediate exchange, 399, 401,
 754–755. *See also* Chemical
 exchange

- Intermediate frequency (IF), 135
Intermolecular chemical exchange, 394, 401, 753–754, 755f
Intermolecular interactions, 753–760
Internal mobility. *See* Dynamics, internal
Interresidue NOE, 788–790, 789f–790f
Intramolecular dynamics. *See* Dynamics, internal
Intramolecular chemical exchange, 393, 401–402
Intrinsic angular momentum, 29–30
Intrinsic amide-solvent exchange rate, 224f
Inverse detection, 535
Inverse Fourier transformation, 137.
See also Fourier transformation
Inversion operator, 52–53, 66
Inversion, population, 16, 57
Inversion recovery, 258–259, 342–346, 686, 688f
IPAP HSQC, 670–672
pulse sequence for, 671f
IRS spin system. *See* Three-spin system
IS spin system. *See* Two-spin system
Isochromats, 218, 247
Isolated spin, 6, 8, 15, 58, 93, 181, 337, 392–400
Isolated two-spin approximation, 507
Isotope editing, 763
Isotope filter, 762–769
breakthrough peaks, 764–765
double difference, 547
double half-filtered NOESY, 765, 766f, 766t, 767
first-order J-filter, 767
half-filter, 763–764
adiabatic, 768
composite, 768–769
gradient, 767, 768
spin-lock, 768
heteronuclear correlation, in, 546–547, 584
second-order J-filter, 764, 768
signal selection by, 763
signal suppression by, 763
third-order J-filter, 768–769
 ω_1 half-filtered NOESY, 765
 ω_2 half-filtered NOESY, 765
Isotope shifts, 732–733, 732t
Isotopic labeling, 533, 727
Isotopomers, 697, 733, 743, 748
Isotropic chemical shift, 22. *See also* Anisotropic chemical shift
dihedral angle restraints from, 804
perturbation mapping of, 756–757
Isotropic mixing, 287–289. *See also* decoupling; TOCSY
coherence transfer efficiency and, 492f, 495f, 611f
Hartmann-Hahn condition and, 288
RDCs and, 291–292
relaxation in, 364
suppression of ROE peaks, 491, 493
Isotropic mixing sequences, 287–288, 291, 490–493. *See also* individual mixing sequences
Isotropic rotor spectral density function, 366
IUPAC-IUBMB-IUPAB Inter-Union Task Group on the Standardization of Data Bases of Protein and Nucleic Acid Structures Determined by NMR Spectroscopy, 262
- J**
- J coupling. *See* scalar coupling
J filter. *See* Isotope filter
JR.NOESY. *See* Jump-return NOESY
Jump-return, 224–226, 233f, 235f
excitation profiles of, 227f
Hahn echo experiment, in, 261f
Jump-return NOESY (JR.NOESY), 511–514
pulse sequence for, 512f
ubiquitin, of, 513
- K**
- Kaiser apodization function, 144–145, 146f, 148, 149f, 150f
Karplus equation, 798
curves, 800f, 802f, 803f
parameterization of, 799, 800t, 801t, 802t, 803t
Ket, 37–38, 47
Kinetics. *See* Chemical kinetics

Kramers-Krönig relations, 143, 357
Kronecker delta, 32

L

Labels, isotopic, 526, 727, 761–763

Labile proton. *See* Amide proton exchange

Laboratory reference frame, 4, 7–8, 102, 104, 106, 107, 108f, 364f, 368, 387

Larmor frequency, 8. *See also* resonance frequency

Lattice, 40, 55, 335

Lattice jump model, 368

Leakage rate constant, 342

Levinson-Durbin linear prediction, 161

Lifetime, 2, 336, 368, 392, 754

Ligand. *See* Protein-ligand binding

Like spins, 360–363, 382

Linear prediction, 160, 161, 234

4D NMR, in, 599–600, 600t

HSQC, of, 162f, 163f

mirror image, 161, 162f, 163f

Linear prediction singular value decomposition (LPSVD), 161

Line-broadening. *See* Apodization; Linewidth

Lineshape. *See also* Apodization; Linewidth

absorptive antiphase lineshape, 413f, 421, 423f, 424–426, 424f, 425f, 470f

absorptive lineshape, 18, 139–140, 140f, 317, 320f

analysis, 684–685, 704–706

chemical exchange, 400f, 507–506

constant time evolution in, 543–546, 545f

dispersive lineshape, 18, 139–140, 140f, 317

dispersive antiphase lineshape, 425f, 470f, 471, 474, 476–477

doubly absorptive, 319, 469

doubly antiphase absorptive, 415, 470, 483

doubly antiphase dispersive, 469–471, 477

doubly dispersive, 319

full-width-at-half-height (FWHH), 18, 139–140, 147–148, 408, 703

Gaussian, 148, 149f, 150f

homogeneous, 18, 544

Lorentzian, 18, 139, 140, 140f, 155, 317

inhomogeneous, 18, 544, 545f

in-phase, 88–90

off-resonance, 168f

phase-twisted, 319, 320f

shimming and, 242

Linewidth

chemical exchange, 400f, 755f

Gaussian, 148

¹H NMR, 406, 408, 407f

HMQC, in, 539–540

HSQC, in, 542, 557

Lorentzian, 18–21, 20f, 139–140, 140f, 729f, 730f

measurement of, 259, 261f

TROSY, in, 387, 571–572

ubiquitin, for, 21, 259, 261f, 408

Liouville operator space, 78–80

Liouville von Neumann equation, 41–43, 79, 351, 356–357

Lipari-Szabo model free formalism, 368–369

Lock, field-frequency, 115f, 116–117
adjustment of, 241
shimming and, 241

Longitudinal relaxation. *See* spin-lattice relaxation

Longitudinal two-spin order, 314, 519, 574, 699, 706, 720

Lorentzian lineshape, 18, 139, 140, 140f, 155, 317

Lorentzian spectral density function, 367, 367f

Lorentzian-Gaussian transformation, 496–497

Lowering operator, 52, 72, 81. *See also* Shift operator

Lower-value (LV), 773, 775

Low-pass filter, 232–234, 233f

LPSVD (Linear prediction singular value decomposition), 161

M

- Magnet, superconducting, 114–117, 116f, 238
- Magnetic equivalence, 353, 361, 412, 483
- Magnetic field. *See also* rf field
- amplitude of, 8
 - anisotropic, 383, 408
 - effective, 8, 10, 16
 - homogeneity of, 115–117, 223, 233, 238, 241–244
 - imaging of, 244, 252
 - inhomogeneity of, 18, 117, 172, 239, 242, 245, 247–251, 293, 311, 408, 703
 - local, 22, 199, 247, 335–336, 367, 383, 387
 - measure of, 8
 - resolution, dependence on, 116
 - secondary, 22
 - shimming, 117, 223, 238–252
 - spherical harmonic expansion of, 239–240
 - static, 4–5, 8, 10, 18, 116, 169, 238
 - stochastic, 336, 345, 367
 - strength of, 4, 8, 10, 116
 - vector, 4, 102
- Magnetic moment, 3–5
- Bloch model and, 7
 - equation of motion, 7
 - expectation value of, 35–37
 - free precession, 14
 - Larmor frequency and, 8, 37
 - precession, 8–9
- Magnetic quantum number, 3, 24–25, 59. *See also* Coherence order; Selection rule
- coherence and, 74–75, 293
 - product basis, in, 59–62
 - spin-1/2 nuclei, for, 24–25
 - spin quantum number and, 2
- Magnetic susceptibility, anisotropic, 110, 666–667
- Magnetization. *See also* Bulk magnetization; Transverse magnetization
- adiabatic pulse, during, 194, 195f
 - average, 399
 - complex, 56, 70, 347
 - equilibrium, bulk, 5, 8–9, 11
 - longitudinal, 5, 12–13, 257, 293–294
 - observable, 55, 68–69
 - sense of precession, 58
 - steady-state, 389, 686
- Magnetization transfer, 280–292. *See also* in-phase coherence transfer; NOESY; ROESY; TOCSY
- Magnetogyric ratio, 3, 3t, 8, 97, 119, 122, 371, 405, 534, 728
- Magnitude mode, 246f
- Main chain directed (MCD) assignment, 786f, 790–792
- Master equation, relaxation and, 351–363. *See also* Relaxation theory
- Matching. *See* Tuning and matching
- Matrix
- density operator representation with, 39, 41, 62
 - direct product, 58–61
 - exponential, 43, 52, 69, 341, 504, 518
 - rotation operator representation with, 52–54, 65–67
 - spin operator representation with, 46–50
 - trace, 39–40
 - unitary, 44, 341
- Matrix representation, 39, 41, 46, 65
- basic kets and, 46
 - exponential operator, of, 43, 69, 341
 - pulse operator, of, 66
 - rotation operator, of, 52–54, 65–67
 - two-spin system, in, 59
- Maximum entropy reconstruction, 159–165, 655, 769–770
- 4D NMR and, 599–600
 - homonuclear 3D, 527
 - HSQC and, 165f
- McConnell equations, 393, 706. *See also* Chemical exchange
- MCD (Main chain directed) assignment, 786f, 790–792
- Medium range NOE, 790f, 798, 807
- Methanol, temperature calibration and, 236
- Methyl group
- assigning, 751

- Methyl group (*Continued*)
- coherence transfer for, 556, 606
 - dynamics, 697
 - fingerprint region for, 418, 419f, 420f, 421f, 453f
 - forbidden cross-peaks and, 453
 - interference in, 360
 - isotopomers, 733
 - linewidths for, 406
 - magnetic equivalence and, 361
 - relaxation and, 259, 361, 369, 453, 483, 696, 700f
 - rotation and, 369, 406, 810
 - selectively protonated, 749–752, 759
 - TROSY, 749, 770
 - ubiquitin, in, 408
- Methylene group
- 2Q spectrum and, 478
 - 3Q spectrum and, 485
 - 3QF-COSY and, 452–453
 - coherence transfer for, 556f, 606
 - DR.COSY and, 436
 - E.COSY and, 456–457
 - interference in, 360
 - isotopomers of, 733
 - Karplus curves for, 801–802, 803f
 - NOESY and, 522, 797
 - R.COSY and, 433
 - ROESY and, 522
 - sensitivity and, 535, 741
 - spin diffusion in, 522
 - stereospecific assignment of, 502, 522
 - TOCSY and, 498, 502
- Methyl-TROSY, 749, 770
- Metric matrix distance geometry, 808
- Microsecond-millisecond time scale, 680, 702
- Microsecond-second dynamics, 702–721
- Millisecond time scale, 702
- Millisecond-second time scale, 702
- Mixed state, 40
- Mixing coefficients, 287, 488, 490, 610
- Mixing period, 273–280, 324, 328f, 330f, 525, 680. *See also* Mixing time
- 2Q and, 471
 - NOESY and, 499, 506, 515, 593
 - R.COSY and, 432
 - ROESY and, 520
 - TOCSY and, 486, 490, 491, 494–496, 499
- Mixing sequences, 490–493. *See also*
- Isotropic mixing sequences
 - chemical exchange and, 481, 493, 522
 - CITY and, 493
 - DIPSI and, 490–493
 - FLOPSY and, 490–492
 - Hartmann-Hahn, 280, 286–288, 492f
 - MLEV and, 490–492
 - NOE and, 491, 493
 - ROE and, 491, 493
 - WALTZ and, 490–492
 - WURST and, 491–493
- Mixing time, 273, 284, 289, 344. *See also*
- Mixing period; *individual NMR experiments*
- MLEV, 206, 208–209, 491, 492f
- mixing sequences and, 490–492
- MLEV-17, 490–491, 492f
- Molecular dynamics, internal.
- See* Dynamics, internal
- Molecular dynamics, restrained (rMD), 808–810
- Molecular reference frame, 367–368
- MPF, 209, 210f, 215, 215f
- MQ. *See* *pQ* NMR
- MQF (Multiple-quantum filtered) 460
- MQF-COSY. *See* *pQF*-COSY
- Multidimensional NMR. *See also*
- individual NMR experiments*
- 2D (Two-dimensional), 273–280
 - 3D NMR, compared with, 327–328, 328f, 329f
 - 3D (Three-dimensional), 327–328, 525–526, 581
 - 2D NMR, compared with, 327–328, 328f, 329f, 525
 - 4D NMR, compared with, 329, 613–614
 - 4D (Four-dimensional), 329–330, 330f
 - 3D NMR, compared with, 329, 613–614
 - sensitivity in, 326–327
- Multiexponential relaxation, 341, 453, 483, 681, 714–715
- Multiple conformations, 799
- Multiple pulse. *See* Composite pulse; Decoupling

Multiple quantum coherence (MQ),
91–92, 473
Multiple quantum echo, 221
Multiple quantum filtered COSY. *See*
*p*QF-COSY
Multiple quantum filtration. *See*
p-quantum filtration
Multiple-quantum (MQ) NMR, 90, 147,
279, 463–486, 465f, 536
chemical exchange in, 715–718
HMQC and, 536, 540
HSQC and, 541, 561
operators and, 90–92, 255, 288
scalar coupling and, 91–92
Multiple-quantum relaxation, in
ubiquitin, 717f
Multiple-quantum transition, 76f
Multiple-spin echo, 221
Multiplet, 23–27, 76–77, 98, 201, 271, 284,
380, 401–402, 455, 535, 548, 571,
572f, 684. *See also* Cross-peaks, fine
structure; Scalar coupling
Lorentzian, 425
Multiplicity, 555
Multiplicity filter, 562
Mutual resolved coupling, 452

N

¹⁵N relaxation rate constants, ubiquitin,
of, 694f, 695f, 705f, 710f, 717f
Natural abundance, 3t, 405, 568, 763,
765, 767
Net magnetization transfer. *See*
In-phase coherence transfer;
Magnetization transfer
NMR active nuclei, 2, 3t
NMR chemical shift time scale, 392,
399–400, 753–754
NMR spectrometer, 114. *See also* Probe;
Shim system
components of, 115f
computer in, 123
data acquisition by, 124
digital filters in, 126–131
lock, field frequency, in, 115f,
116–117
oversampling in, 127–131
quadrature detection in, 132–136,
133f, 134f

receiver in, 115f, 123
rf transmitter in, 115f, 120–122
NOE (Nuclear Overhauser effect), 680.
See also Cross-relaxation; NOESY;
ROE
accuracy of, 798
calibration of, 510, 597–598
classifications
intraresidue, 787f, 788
long range, 786f, 806–807
medium range, 790f, 797–798,
806–807
sequential, 786f, 788–790, 789f,
790f, 806–807
coverage, 798
deuteration and, 745, 748–749
difference, 389–390
distance dependence of, 797–798
enhancement, 345, 389–390
global fold and, 807f
heteronuclear, 290, 694f
mixing sequences and, 491, 493
protein-ligand binding and, 761–762
rate constant, 390–391, 796–797
ROESY, contribution to, 520
steady-state, 389–390, 690, 700,
812
structural restraints and, 510, 797–798
transferred, 761
transient, 289, 344–345, 390–391, 700,
805
ubiquitin for, 694f, 789f, 790f
NOESY (Nuclear Overhauser Effect
Spectroscopy), 289–290, 502–517.
See also Cross-relaxation; Isotope
filter; Jump-return NOESY;
ROESY; ZZ-exchange; *individual*
NMR experiments
baseline in, 506, 509
buildup series for, 510
chemical exchange and, 397, 511,
522
deuteration and, 745, 748–749
double half-filtered, 765, 766f, 766t,
767
excitation sculpting and, 513–514,
514f
experimental protocol for, 506–510
Hahn echo and, 502–503, 503f
initial rate regime for, 507

- NOESY (Nuclear Overhauser Effect Spectroscopy) (*Continued*)
 mixing period in, 502, 506–508, 510, 515
 phase cycling in, 509, 511–512
 presaturation in, 506, 511, 513f
 processing of, 510
 product operator analysis for, 503–506
 protein-ligand binding and, 761–762
 pulse sequence for, 503f, 504f, 512f, 515f
 recycle delay in, 506, 514
 relayed, 515–517
 pulse sequence for, 515f
 ubiquitin, of, 516f
 resonance assignment and, 786, 788–792, 789f, 790f
 spin diffusion in, 506–508
 ubiquitin, of, 509f, 513f, 514f, 516f
 variants, of, 511–517
 ω_1 half-filtered, 765
 ω_2 half-filtered, 765
 zero-quantum peaks in, 490, 505, 508–509
 z-filtration in, 490, 508–509
 ZQ peaks in, 490, 505, 508–509
 NOESY-HSQC, 582–589. *See also*
 HSQC-NOESY; HMQC-NOESY-
 HMQC; HMQC-NOESY-HSQC;
 HSQC-NOESY-HSQC
 pulse sequence for, 584f
 product operator analysis for, 582–584
 schematic of, 586f
 ubiquitin for, 587f, 589f
 NOESY-NOESY, 528–529
 NOESY-TOCSY, 525–529, 526f
 Noise, 119, 123, 126–129, 131–132, 158, 163, 253. *See also* Signal-to-noise ratio
 Noise, t_1 , 262, 453
 Nonadiabatic relaxation processes, 336–337, 349, 350
 Nonlinear nonresonant phase shift. *See* Phase shift, nonresonant
 Nonresonant frequency shift, 170–172
 Nonresonant phase shift. *See* Phase shift, nonresonant
 Nonsecular, 355, 361–362, 380, 399, 684
 Nonselective inversion recovery, 258–259, 260f, 344, 346f
 Normalization, 30, 49, 79
 N/P selection, 316–317, 322, 576. *See also* Frequency discrimination
 N-type signal, 316–317, 319, 322
 Nuclear magnetic moment, 3–6
 Nuclear magnetism, 2, 5–6
 angular momentum and, 3
 magnetogyric ratio and, 3
 nuclear spin states and, 5–6
 Nuclear Overhauser effect. *See* NOE
 Nuclear Overhauser effect (NOE)
 classifications
 intraresidue, 787f, 788
 long range, 786f, 806–807
 medium range, 790f, 797–798, 806–807
 sequential, 786f, 788–790, 789f, 790f, 806–807
 Nuclear Overhauser Effect Spectroscopy. *See* NOESY
 Nuclear shielding, 22, 51, 102, 107
 Nuclear spin angular momentum. *See* Angular momentum
 Nuclei
 active, 2, 3t
 properties of, 3t
 quadrupolar, 2
 Nutation axis, 120–122
 Nyquist frequency, 124–126, 126f
- O**
- Observable
 operator, 32, 55–56, 89–90, 300, 308
 transition, 6, 25
 Observable quantity, 31–34, 41
 Observation operator, 55–56, 300, 308
 Off-resonance
 180° (π) pulse, 167f, 169, 178f
 90° ($\pi/2$) pulse, 166–169, 167f, 168f
 phase shift, 168–171
 pulse, 165–169
 Offset
 baseline, 155–157, 157f, 307–308, 326
 resonance, 10, 58, 121
 ω_1 half-filter. *See* Isotope filter
 One-dimensional NMR. *See* 1D ^1H NMR

- One-pulse acquire experiment, 16–18, 19f, 56–58, 263, 265
 recycle delay, for, 257–258, 258f
 signal-to-noise of, 158–159
 two-spin system, for, 68–70
 ubiquitin, of, 150f, 233f, 235f, 267f, 272f
- One-spin operators, 46, 70
- On-resonance
 pulses, 16, 17f, 51, 85–86
 rf field, 11, 12f
- Operator. *See also* Angular momentum operator; Basis operator; Cartesian operator; Density operator; Hamiltonian; Product operator; Tensor Operator; Trace
 adjoint, 32
 bilinear, 91, 287–288
 compatible, 34
 commuting, 34
 direct product, 60–61
 equilibrium density, 55–56, 75, 294
 expansion of, 78–79
 exponential, 42–43, 45–46
 Hermitian, 31–32, 39, 41
 identity, 38, 55–56
 inversion, 52–53, 66
 matrix representation, 39, 41, 43, 46–47
 MQ, 90–92
 observable, 32, 55–56, 89–90, 300, 30
 observation, 55–56, 300, 308
 one-spin, 46, 70
 orthogonal, 79
 Pauli spin, 46–50
 projection, 39, 74
 rotation, 50–54, 65–67, 84, 85f
 shift, 70, 72, 73t, 75, 80–81, 90, 293, 316, 356
 similarity transformation of, 44–45, 48, 341
 single element, 80–83, 99, 353, 450, 658
 single transition shift, 81, 83f, 100, 294, 460
 space, 78–80
 spherical tensor, 103–104, 353
 two-spin, 70, 71t, 73t
 unitary, 44
- Optimal control theory, 179, 726
- Order parameter
 Saupe, 109
 Lipari-Szabo, 368–370
 ubiquitin, for, 812f
- Orientational correlation function.
See Stochastic correlation function
- Orientational spectral density function.
See Spectral density function
- Orthogonal operator, 79
- Orthogonality, 32, 79. *See also* Normalization
- Orthonormal set, 32–33, 38
- Orthonormality, 32
- Oscillator, damped, 139. *See also* FID; Interferogram
- Oscilloscope, 237, 238f, 253
- Out-and-back coherence transfer, 613
- Out-and-stay coherence transfer, 632
- Oversampling, 126–129, 266
- ## P
- Paramagnetic relaxation, 365, 370
- Paramagnetic effects, 797
- Parseval's theorem, 138
- Parts per million (ppm), 23, 262
- PAS. *See* Principal axis system
- Pascal triangle, 412
- Passive scalar coupling. *See* Scalar coupling
- Pauli spin matrices, 37, 46
- P.COSY (Purged Correlation Spectroscopy), 427–428
- Peaks. *See also* Cross-peaks; Direct peaks; Remote peaks
 back-transfer, 527
 breakthrough peaks, J-filter, 764
 diagonal, 277, 290, 452–453, 475, 593, 597
 zero-quantum, 490, 505–506, 508–509
 suppression of, 314, 508–509
- Peakshape. *See* Lineshape
- Peak-to-peak voltage, 253
- P.E.COSY (primitive E.COSY), 427
- PEP (Preservation of equivalent pathways), 500–502, 560–572. *See also individual NMR experiments*
- ¹³C heteronuclear correlation, in, 598

- PEP (Preservation of equivalent pathways) (*Continued*)
- ¹⁵N heteronuclear correlation, in, 554f, 560–566, 570f
 - gradient-enhanced, 574, 575–578, 577f
 - HSQC-TOCSY, in, 593
 - triple-resonance, in, 620f, 627–628
 - TROSY, in, 566–570, 570f, 620f, 628
 - relaxation, 562–563, 568–570
 - sensitivity enhancement, 562–563
 - spin multiplicity in, 560–561
 - TOCSY, in, 500–502, 593
- Peptide
- chemical shifts, random coil, 23, 408, 782, 794, 804
 - ¹H 1D spectrum of, 272f
 - scalar coupling in, 799
 - test sample, as, 255–256
- Perdeuteration, 727
- ¹³C α decoupling and, 734
 - isotope shifts and, 732–733, 732t
 - relaxation and, 728–729, 729f, 734
 - sensitivity and, 729–732
- PFG. *See* Pulsed field gradient
- Phase angle, 10, 37, 120, 166, 167f, 720
- Phase correction
- aliasing and folding in, 125–126
 - baseline distortion, from, 156–158, 156f, 157f, 324
 - first-order, 131, 152, 154f, 155
 - nonresonant phase shift for, 170–172, 621–622, 645
 - pivot, 152, 155
 - zero-order, 152, 153f, 154f, 155
- Phase cycle, 292–307. *See also*
- Coherence transfer pathway;
 - Frequency discrimination; Isotope filter; Quadrature detection; *individual NMR experiments*
 - artifact suppression in, 307–310
 - axial peak suppression, 309–310, 323–324
 - cogwheel, 307
 - CYCLOPS (Cyclically ordered phase sequence), 266, 307–308
 - EXORCYCLE, 266, 309, 313
 - first pulse, for, 306–307
 - grouping pulses in, 306–307
 - heteronuclear correlation, for, 546–548
 - last pulse, for, 307
 - limitations of, 310–311
 - p*-quantum, 465, 465f
 - p*-quantum filtration, 437f, 438, 461t
 - repetition rate artifacts, 258, 310–311, 444–445
 - saving time in, 305–307, 310–311
 - selectivity of, 305
- Phase error, 123, 127, 154–156, 170, 266, 539, 622, 645, 659
- Phase map, 246–249, 250
- Phase modulated signal, 316–317
- Phase, NMR spectrum of, 139, 151–158
- baseline distortions, 156–158, 156f, 157f
 - frequency-dependent, 151–152, 154f
 - frequency-independent, 151–152, 153f, 154f
 - initial, 139
 - sampling delay and, 154–156, 156f
- Phase, pulse, 10–11, 168–171
- Phase sensitive. *See* Frequency discrimination; Quadrature detection
- Phase shift, resonance
- nonresonant, 9, 170–172
 - cosine modulation, compensatory, 172
 - delays, compensatory, 597, 599
 - phase shift compensatory, 171–172, 621–622, 645
 - pulses, compensatory, 172, 579–580
 - off-resonant pulse, 166–169, 167f, 168f
 - receiver phase by, 295–300, 302–306
 - rf phase by, 298
- Phase, spatially dependent, 218, 220, 311, 576
- Phase-incremented pulse (PIP). *See* Phase-modulated pulse
- Phase-modulated pulse, 181–189, 185f
- vector representation of, 190f
- Phase-twisted lineshape, 319, 320f
- Phasing, 151–156, 153f, 154f, 156f. *See also* Phase correction
- in multidimensional NMR, 323–326

- Picosecond-nanosecond dynamics, 685–701
- ^{15}N backbone amide, 686–692, 688f, 693f–695f
- $^{13}\text{CH}_2\text{D}$ methyl, 693, 696–699, 696f, 698f, 700f
- ^{13}CO backbone, 699–700, 701f
- PIP (Phase-incremented pulse). *See* Phase-modulated pulse
- Pivot, 152, 155
- Planck's law, 6
- Polar angle, 110, 365, 368, 804–805
- Polarization transfer. *See* Coherence transfer
- Poles, signal, 160
- Polyacrylamide gels, RDC and, 668–669
- Population. *See also* Spin-lattice relaxation
- Boltzmann distribution, 5, 55–56, 335
- density matrix, 55–56
- effects of rf pulse on, 75
- equilibrium, 5
- inversion, 16, 57
- Solomon equation for, 339–340
- transfer, 98–99, 397f, 706
- Population transfer, chemical exchange and, 397f, 706
- Post-acquisition signal processing, 232–234
- Post-acquisition solvent suppression, 232–234
- Power, 253–254
- Power spectral density function. *See* Spectral density function
- p*Q NMR (Multiple quantum NMR), 463–465, 465f. *See also* 2Q NMR; 3Q NMR
- p*QF COSY (Multiple quantum filtered COSY), 437–440, 437f. *See also* 2QF-COSY; 3QF-COSY
- p*-quantum filtration, 437–440. *See also* 2QF-COSY; 3QF-COSY
- PR. *See* Projection-reconstruction
- Preacquisition delay, 265–266, 494
- Preamplifier, 115f, 119, 123, 133f, 253
- Precession. *See* Free precession
- Precision, three-dimensional structures of, 798, 805–806, 809–810
- Prediction order, 160
- Preparation period, 271, 273f, 279–280, 328f, 330f, 680
- Presaturation, 223–224, 233f
- NOESY and, 513f
- Preservation of equivalent pathways. *See* PEP
- Pre-TOCSY, 428–429, 428f, 499
- Pre-TOCSY COSY, 428–429, 428f, 499
- Primitive Exclusive Correlation Spectroscopy (P.E.COSY), 427
- Principal axis system (PAS), 22, 103–104, 108f, 110
- Principal component of tensor, 22, 103, 110, 110
- Principal value of tensor, 22, 103, 110, 110
- PR-NMR (Projection reconstruction-NMR), 773–775, 774f
- Probability density, 30, 40, 107, 109, 111, 350
- Probability, transition, 334–335
- Probe, 114–120, 115f, 118f. *See also* rf coil
- actively shielded gradient, 217, 219–220
- cryogenic, 119
- impedance, 118–119, 237
- quality factor, 118, 159
- resonance frequency, 237
- tuning and matching, 119, 237–238, 238f
- PROCHECK, 809–810
- PROCHECK-NMR, 809–810
- Prochiral groups, assignment of, 502, 798
- Product basis, two-spin, 24, 58–61, 62–64
- Product operator. *See also* Basis operator; Cartesian operator; Density operator; Hamiltonian; Operator; Tensor Operator; Trace
- Cartesian basis, in, 58–61, 70, 71t, 80
- shift, 70, 72, 73t, 75, 80–81, 90, 293, 316, 356
- single element, 80–83, 99, 353, 450, 658
- single transition shift, 81, 83f, 100, 294, 460
- transformation of, 84–88, 85f, 87f

- Product operator formalism, 77–102
 evolution of, 84–88, 85f, 87f
 cascades during, 86, 88
 chemical shift, 84, 85f
 commuting operators of, 88
 double quantum (DQ) coherence of, 90–93
 free precession, 84–86, 87f, 90–92
 rotation, 84, 85f
 scalar coupling, 84–85, 87f, 92
 zero quantum (ZQ) coherence of, 90–93
 pulses, 85–86, 87f
 relaxation, 337–338, 357, 359, 362
 transformation, 84–88, 85f, 87f
 Profile edge shimming, 252
 Progressively connected transition, 77, 460
 Projection operator, 39, 74
 Projection-reconstruction (PR), 773–775, 774f
 Propagator, 45–46, 54, 79–80, 84, 337–338
 Protein. *See also* Ubiquitin; calbindin D_{28k}
 chemical shifts for, 784f, 785f, 795f, 796f
 correlation time for, 18, 21
 denatured, 407f, 408–409
 hydration layer, 21
 hydrodynamic radius, 21
 linewidth, 20f, 729f, 730f
 scalar coupling constants for, 800f, 800t, 801t, 802f, 802t, 803f, 803t
 specific volume, 21
 Protein structure. *See* Structure determination
 Protein-ligand binding, 753–769. *See also* Isotope filter
 amide proton exchange, 760
 chemical exchange regime, 753–755
 chemical shift mapping, 756–757
 cross saturation, 757–759
 NOESY and, 761–769
 relaxation for detecting, 759–760
 Proton decoupling, 552–553, 555–557, 570f, 626–627
 Proton detection, 535
 Proton exchange. *See* Amide proton exchange
 Protonation, 749–753
 amide $^1H^N$ spins, 727–728
 aromatic rings, 752–753
 ILV, 751
 methyl-selective, 749–752
 ubiquitin, ILV NOESY, of, 752f
 Pseudo-diagonal, 471, 475f, 597
 Pseudo-energy, from restraints, 808–810
 Pseudo-first-order rate constant, 394
 P-type signal, 316–317, 319, 322
 Pulse, 14–15. *See also* 180° pulse; 90° pulse; Adiabatic pulse; Composite pulse; Phase-modulated pulse; Pulsed field gradient; Selective pulse; Shaped pulse
 broadband inversion (BIP), 179
 Cartesian product operators and, 84
 coherence order, effect of, 294
 evolution during, 166–169
 product operator formalism, in, 84–85
 excitation null for, 169–170
 flip angle, 11, 52
 Hamiltonian and, 51
 homospoil, 220, 574
 length, 11, 171, 253–255
 off-resonance, 166–169, 167f, 168f, 169, 178f
 on-resonance, 16, 17f, 51, 85–86
 phase, 10–11, 168–171
 purge, 538, 541, 659, 663
 read, 224, 226, 511
 rf field strength of, 8
 rotation angle, 11, 52
 rotation operator, 50–54, 65–67, 84, 85f
 spin lock purge, 173, 227–230, 228f, 232, 573, 585, 767, 768
 Pulse programmer, 115f, 120
 Pulse sequence, 43
 gradient shimming, 248f
 INEPT, 96–99
 pulse width calibration, for, 254–257, 256f
 out-and-back, 613
 out-and-stay, 632
 spin-state-selective (S^3CT), 99–101
 spin echo, 93–96
 straight-through, 632

- Pulse sequence, specific experiment
 - heteronuclear
 - CE-TROSY, 673f
 - Cross-saturation, 758f
 - Double half-filtered NOESY, 766f
 - Fast HSQC, 575f
 - ^1H - ^{13}C CT-HSQC, 579f
 - HCCH-COSY, 603f
 - HCCH-TOCSY, 610f
 - HMQC, 537f
 - HMQC-NOESY-HMQC, 594f, 596f, 598f
 - HNHA, 662f
 - HSQC, 537f, 554f, 575f, 577f, 579f
 - HSQC-NOESY, 592f
 - HSQC-NOESY-HSQC, 746f, 750f
 - HSQC-TOCSY, 591–593
 - IPAP HSQC, 671f
 - PEP-HSQC, 554f
 - PFG-PEP-HSQC, 577f
 - PFG-TROSY, 577f
 - NOESY-HSQC, 584f
 - TOCSY-HSQC, 590f
 - TROSY, 554f, 575f, 577f, 673f
 - homonuclear
 - 2Q, 465f
 - 2QF-COSY, 437f, 439f
 - 3Q, 465f
 - 3QF-COSY, 437f
 - COSY, 410f
 - COSY- β , 426–427
 - DR.COSY, 435f
 - E.COSY, 457, 461t
 - JR.NOESY, 512f
 - NOESY, 503f, 504f, 512f, 515f
 - NOESY-TOCSY, 526f
 - P.COSY, 427–428
 - p*Q NMR, 464–465
 - p*QF-COSY, 437–439
 - pre-TOCSY COSY, 428–429, 499
 - R.COSY, 430f
 - relayed NOESY, 515f
 - ROESY, 521f, 524f
 - SCUBA, 429
 - TOCSY, 488f, 489f
 - 1D ^1H
 - 1-3-3-1, 224–227
 - excitation sculpting, 228f
 - Hahn echo, 228f, 229f, 265–266, 704f
 - jump-return, 224–225, 261f
 - one-pulse acquire, 263–264
 - WATERGATE, 229–230
 - water flip-back, 229f
- Relaxation measurement
 - 1D inversion-recovery, 258
 - $^{13}\text{CH}_2\text{D}$, for, 696f, 698f
 - ^{13}CO , for, 701f
 - CPMG, 712f, 715f
 - Hahn spin echo, 261f, 704f, 719f
 - multiple quantum, 717f
 - ^{15}N , for, 688f, 695f, 704f, 709f, 717f
 - $\text{R}_{1\rho}$, 709f
 - relaxation interference, 695f
 - TROSY-CPMG, 721f
 - TROSY Hahn spin echo, 719f
 - ZZ-exchange, 707f
- Triple-resonance
 - CBCA(CO)NH, 643f
 - CBCANH, 648f
 - CC(CO)NH-TOCSY, 742f
 - HBHA(CBCACO)NH, 645
 - HBHA(CBCA)NH, 649
 - HCACO, 637
 - (HCA)CONH, 641
 - H(CA)NH, 633f
 - HNCA, 619f–620f, 736f
 - HNCACB, 651f, 738f
 - HN(CACB)CG, 740
 - HN(CA)CO, 640f
 - HNCA-*J*, 657f
 - HNCO, 638f
 - HN(CO)CA, 630f
- Pulse width calibration, 252–257
 - pulse sequences for, 256f
- Pulse-acquire. *See* One-pulse acquire experiment
- Pulsed field gradient probe, actively shielded, 217, 220, 246f, 251f
- Pulsed field gradient (PFG), 217–221, 311–315
 - artifact suppression, 313–313, 577f
 - coherence selection, 311–314, 574–578
 - dephasing by, 219f
 - frequency discrimination by, 322–323, 574–578
 - limitations of, 314–315
 - magic-angle, 221, 444

Pulsed field gradient (*Continued*)
 radiation damping and, 219, 231–232, 440, 514, 587
 recovery delay for, 315
 shape factor for, 311
 solvent suppression using, 229f, 230–232, 573–574
 strength of, 218, 221
 three-axis, 220, 444
 z-axis, 217–218
 Pulse-interrupted free precession, 280–284
 Pure state, 40, 74
 Purged Correlation Spectroscopy (P.COSY), 427–428
 Purity, sample, 409, 418

Q

Quadrature detection, 117, 132–136, 133f, 134f, 321. *See also* Frequency discrimination; States; Time-proportional-phase-incrementation; TPPI-States
 Quadrature glitch, 307–308
 Quadrature images, 135–136, 264–265, 307–308
 Quadrature pair, 132, 564
 Quadrupolar Hamiltonian, 680
 spatial functions for, 366t
 spherical harmonic functions for, 350t
 tensor operators and, 385t
 Quadrupolar nuclei, 2, 384
 Quadrupolar relaxation, 370–371, 383–384, 385t
 techniques for, 693, 696–699, 696f, 698f
 ubiquitin, of, 700f
 Quadrupole moment, 2, 384. *See also* Electric field gradient tensor
 axial symmetry of, 366
 principal axis of, 365
 spatial function for, 366
 spherical harmonic functions for, 350t
 tensor operators and, 385t
 Quality factor, 118, 159, 211
 Quantitative *J*-correlation, 656, 660, 662, 670

Quantum mechanical relaxation theory, 351
 Quantum mechanics
 density matrix, 37–40
 multispin systems, of, 58–70
 postulates of, 29–37
 Quantum number, angular momentum, 2. *See also* Magnetic quantum numbers
 Quantum statistical mechanics, 40–41
 Quaternion formalism, 15, 177

R

R_1 . *See* Spin-lattice relaxation
 $R_{1\rho}$, 382. *See also* Rotating frame relaxation
 R_2 . *See* Spin-spin relaxation
 R_2^* , 18, 408, 544
 Radiation damping, 123, 222, 232, 241, 248–249, 440, 511–512, 514, 574, 586–587
 Radiofrequency field. *See* rf field
 Raising operator. *See* Shift operator
 Rance-Kay processing, 563, 564, 567–568
 Random coil chemical shifts, 23, 408, 782, 794, 804
 Random field. *See* Stochastic field
 Random fractional deuteration, 727
 isotopomers in, 733
 relaxation and, 728–729, 730f, 741
 Random function. *See* Stochastic function
 R.COSY, 429–436
 experimental protocol for, 432
 processing of, 432
 product operator analysis for, 430–431
 pulse sequence for, 730f
 ubiquitin, of 434f
 RCT. *See* R. COSY
 RD (Reduced dimensionality), 771–772
 RDC. *See* Residual dipolar coupling
 RE-BURP, 182f, 184f, 186
 Read pulse, 224, 511, 528
 Real Fourier transformation, 141, 317, 318f, 319, 321–322. *See also* Fourier transformation

- Receiver, 115f, 123
 gating delay, 157–158, 266
 phase shift, 297–298
- Recovery delay, pulsed field gradient, 315
- Rectangle function, 144–145
- Recycle delay, 98, 257–258, 258f
 double diagonal artifacts and, 444–445, 445f
 Ernst angle and, 257, 258f
 NOESY, for, 506
 phase cycling and, 306
 signal-to-noise and, 158, 534
- Redfield equation, 350–359. *See also* Relaxation theory
- Redfield kite, 358f
- Redfield's method for frequency discrimination, 321
- Reduced dimensionality (RD), 771–772
- Reduced static field, 10
- Reference, chemical shift, 23, 262–263
- Reference frame
 alignment, 108, 804–805
 frequency modulation (FM), 191–192, 196
 interaction, 351–352
 laboratory, 4, 7–8, 102, 104, 106, 107, 108f, 364f, 368, 387
 molecular, 107, 367–368
 rotating, 7, 10–11, 11f, 43–46, 364f
 tilted, 10–11, 11f, 363–364, 364f, 381–383
- Reference frequency, 132, 135, 322
- Refinement, 3D structures of, 805, 809–810
- Refocused INEPT, 98–99, 553, 556f.
 See also INEPT
- Refocusing, by 180° pulses, 93–96
- Regressively connected transition, 77, 460
- Relaxation, 333. *See also individual NMR experiments*
 adiabatic, 336, 337, 349–350
 autorelaxation, 340, 357, 359, 362, 375, 382–383, 692
 average, 364, 380–381, 383, 396, 540, 542, 572, 681–685, 713, 718, 720, 748
 biexponential, 343–345, 681
 coherence, dephasing by, 336
 constants in, 337
 cross-correlation. *See* Relaxation interference
 cross-rate constant, 342
 cross-relaxation. *See* Cross-relaxation
 chemical shift anisotropy (CSA), 22, 349–350, 383–384, 385t, 386–387, 502–506, 571, 680
 decay constant, 139, 399, 400, 692
 deuteration and, 728–729, 729f, 730f
 dipolar, 20f, 359–360, 365–366, 370–383, 377t, 386, 502, 571–572, 680, 685. *See also* Cross-relaxation; NOE; NOESY; ROE; ROESY
 dynamic frequency shift, 356–357
 extreme narrowing, 367, 384, 391
 forbidden, 360, 453, 483
 heteronuclear multiple quantum coherence for, 540
 heteronuclear single quantum coherence for, 378–381, 542
 high temperature limit, 356, 365
 inhomogeneous broadening, 18, 534–544, 545f
 initial rate regime, 343–345, 507, 681, 797
 interference. *See* Relaxation interference
 ¹H relaxation rate constants, 20f, 729f, 730f
 laboratory reference frame, 357–358, 363, 364f, 368, 383, 686–701
 leakage rate constant, 342
 like spins, 360–363, 382
 Lipari-Szabo formalism, 368–369
 longitudinal. *See* Spin-lattice relaxation
 magnetic equivalence in, 353, 360, 391–392
 master equation, 351–359
 monoexponential, 344–345, 683
 multiexponential, 341, 453, 483, 714
 ¹⁵N relaxation rate constants, ubiquitin, of, 694f, 695f, 705f, 710f, 717f
 nonsecular contribution to, 355, 361–362, 380, 399, 684
 paramagnetic, 365, 370

- Relaxation (*Continued*)
- perdeuteration and, 728–729, 729f
 - population, 335–336, 339–340.
 - See also* Spin-lattice relaxation
 - product operator, for, 337–338, 357–358, 359, 362
 - quadrupolar, 365, 370–371, 383–384, 385t
 - random fractional deuteration and, 728–729, 730f, 741
 - rate matrix, 341, 357–358, 358f, 372, 378, 395, 397, 506, 680–684
 - Redfield kite, 358f
 - rotating frame, 363–364, 364f, 381–383, 390–391
 - scalar, 370, 381, 387–388, 401, 406, 734
 - of first kind, 387–388
 - of second kind, 387–388
 - secular approximation in, 358, 360–363, 372, 376, 380, 683
 - self-decoupling by, 381, 401, 660, 799
 - Solomon equations, 338–346, 375, 503
 - relaxation interference and, 386
 - spatial functions for, 366t
 - spin diffusion, 367, 511, 515, 528, 745, 759, 797, 798
 - NOESY and, 506–508, 522
 - ROESY and, 522
 - spin diffusion regime and, 367
 - spin-lattice. *See* Spin-lattice relaxation
 - spin-spin. *See* Spin-spin relaxation
 - superoperator, 356, 361, 363
 - transverse. *See* Spin-spin relaxation
 - ubiquitin
 - order parameters for, 812
 - relaxation decay for, 693f, 710f
 - relaxation dispersion in, 710f
 - relaxation rate constants for, 694f, 695f, 700f, 705f, 717f
 - unlike spins, 360–363, 714
- Relaxation interactions and mechanism. *See also* Chemical exchange
- chemical shift anisotropy (CSA), 22, 349–350, 383–384, 385t, 386–387, 502–506, 571, 680
 - principal axis for, 349–350, 365, 387
 - spatial functions for, 366t
 - spherical harmonic functions for, 350t
 - tensor operators and, 384t
 - dipolar relaxation, 20f, 359–360, 365–366, 370–383, 377t, 386, 502, 571, 680. *See also* Cross-relaxation; NOE; NOESY; ROE; ROESY
 - picosecond-nanosecond dynamics and, 685
 - principal orientation of, 365
 - spatial variables for, 366t
 - spherical harmonic functions for, 350t
 - tensor operators for, 372t, 382
 - TROSY and, 572
 - paramagnetic, 365, 370
 - quadrupolar, 370–371, 383–384, 385t
 - spatial functions for, 366t
 - spherical harmonic functions for, 350t
 - tensor operators and, 385t
 - scalar, 370, 381, 387–388, 401, 406, 734
 - of first kind, 387–388
 - of second kind, 387–388
- Relaxation interference, 359–360, 385–387
- CRIPT and, 726
 - CSA-CSA, 692
 - CSA-dipolar, 359–360, 373, 385–387, 570–572, 672, 686, 692, 695f, 703–704, 713
 - dihedral angle and, 801
 - dipole-dipole, 692
 - forbidden cross-peaks from, 360, 453, 483
 - experimental methods for, 683–685, 692–693
 - multiplet structure and, 684–685
 - spectral density function for, 369–370
 - TROSY and, 570–572, 718–721
- Relaxation measurements
- $^{13}\text{CH}_2\text{D}$, for, 696f, 698f
 - ^{13}CO , for, 699–700, 701f
 - CPMG, 712f, 715f
 - Hahn spin echo, 261f, 704f, 719f
 - inversion recovery, 258–259, 342–346, 686, 688f
 - Multiple quantum, 717f
 - ^{15}N , for, 688f, 695f, 704f, 709f, 717f
 - NOE difference, 388–390

- NOESY. *See* NOESY
 $R_{1\rho}$, 709f
relaxation interference, 695f
ROESY. *See* ROESY
steady-state NOE, 388–390, 690–691, 688f, 699–700, 701f
transferred NOE, 760–761
transient NOE, 344–345, 346f
TROSY-CPMG, 721f
TROSY Hahn spin echo, 719f
ZZ-exchange, 707f
Relaxation mechanisms. *See* Relaxation interactions and mechanisms
Relaxation theory
 Bloch equations, 12–14, 337–338, 346–350
 Bloch-McConnell equations, 393–400
 quantum mechanical, 350–351
 semi-classical (BMR), 350–364, 358f
 Solomon equations, 338–346, 343f, 375, 503–504, 506–507
Relaxation-compensated CPMG, 712f, 713–715, 715f
Relayed coherence transfer (RCT). *See* R.COSY
Relayed COSY. *See* R.COSY
Relayed NOESY, 515–517
 pulse sequence for, 515
 ubiquitin, of, 516
Remote peak
 2Q and, 469–472, 471f, 472t, 475, 475f, 479f
 3Q and, 484–485
Reorientation. *See* Diffusion
Repetition rate artifacts, 258, 310–311, 444–445
Residual dipolar coupling (RDC), 110, 665–673, 810–811
 alignment media for, 667–669
 coherence transfer using, 291
 measuring, 670–673
 restraint, from 751, 804–805
 ubiquitin, for, 669f, 812f
Resistance, 119–120, 253–254
Resolution
 digital, 141, 158
 constant-time evolution, in 544, 549–550
 semi-constant time evolution, in, 691–692
 enhancement, 147–148, 149f, 150f
 magnetic field, dependence on, 23, 116
Resonance assignment
 ^1H , 782–791
 isotopically labeled proteins, for, 792–794
Resonance degeneracy. *See* Chemical shift, degeneracy
Resonance dispersion, 179, 236, 405, 407f, 408, 785, 791
Resonance frequency, 8, 11, 125–132, 237, 324
Resonance lineshape. *See* Lineshape; Linewidth
Resonance linewidth. *See* Lineshape; Linewidth
Resonance offset, 10, 58, 121
Restrained molecular dynamics (rMD), 808–809
Restraint
 covalent, 808
 dihedral angle, 798–804, 800f, 802f, 803f, 804, 806–807
 distance, 510, 522, 751–752, 797–798, 806–807
 hydrogen-bond, 805–806, 807
 NOE, 507–508, 510, 522, 751–752, 797–798, 806–807
 Paramagnetic effects, 797
 protein complexes, for, 756, -769
 RDC, 804–805
 ROE, 522
 residue, per, 798
 scalar coupling, 798–804, 800f, 802f, 803f, 806–807
 violations, 809
rf coil, 116f, 117–119, 118f
rf field
 amplitude of, 8, 9–11, 120
 B_1 , 9, 11f
 Bloch-Siegert shift of, 9, 170
 counter-rotating, 9, 50, 170
 Hamiltonian for, 50
 inhomogeneity of, 172–174, 178f, 228–230, 364, 517, 519
 nonresonant, 9, 170

rf field (*Continued*)
 off-resonance, 10–11, 11f, 12f, 166–167
 on-resonance, 10, 11f, 12f
 phase of, 9–10, 120
 strength of, 8, 9–11, 120
 rf power meter, 237, 238f
 rf transmitter in, 115f, 120–122
 Ridges, 416, 493, 785
 Rigid rotor, 366–367
 Ring flip, aromatic, 369, 480, 810
 Ringing, 265
 rMD (restrained molecular dynamics), 808–809
 RMSD (Root-mean-square deviation), 798, 808–809
 ROE (Rotating frame Overhauser effect), 363–364, 383, 390–391, 491, 493, 502, 517, 520–522. *See also* Cross-relaxation; NOE
 ROESY (Rotating-Frame Overhauser Effect Spectroscopy), 517–524. *See also* Cross-relaxation; NOESY; ROE
 chemical exchange in, 522, 524
 experimental protocol for, 520–521
 mixing time for, 520–521
 vs. NOESY, 517, 522
 product operator analysis for, 517–520
 processing of, 521–522
 pulse sequence for, 521f, 524f
 relaxation in, 363–364, 383, 390–391
 resonance offset in, 518–520
 rf inhomogeneity in, 364, 517
 spin diffusion in, 522
 ubiquitin, of, 523f
 variants of, 524
 Roll, baseline, 157, 266
 Root-mean-square deviation (RMSD), 798, 808–809
 Root-mean-square voltage, 253
 Rotating frame, 7, 10–11, 11f, 43–46, 364f
 Rotating frame Overhauser effect. *See* ROE
 Rotating frame transformation, 43–46, 50–51, 363–364

Rotating-Frame Overhauser Effect Spectroscopy. *See* ROESY
 Rotating-frame relaxation, 363–364, 364f, 381–383, 390–391
 Rotation, convention for, 10, 58, 121–122
 Rotation operator, 50–54, 65–67, 84, 85f
 Rotational correlation time, 20f, 21. *See also* Diffusion
 Rotational diffusion. *See* Diffusion
 R-SNOB, 182f, 184f, 186

S

S³CT. *See* Spin-state selective coherence transfer
 SA (Simulated annealing), 809
 SAIL (Stereo-array isotope labeling), 727
 Sample preparation, 234–236
 Samples, test, 255
 Sampling, 124–125
 decimation in, 129
 nonuniform, 770–771
 Sampling delay, 139, 154, 156, 161, 172, 324
 acquisition, adjusting, 154, 324, 546, 655
 indirect evolution, 324, 770
 adjusting in, 324–326, 546, 658
 compensating with added delay, 597
 Sampling interval, 124–125, 140, 158, 265, 320–321
 Sampling rate, 124–129, 132, 265
 Sampling theorem, 124–126, 126f
 SAR (Structure-activity relationship), 736
 Saturation, 253, 259, 271, 477f, 585, 685, 688f, 689–691, 701f, 755f, 757–759
 Saturation transfer, 223–224, 226, 228, 232, 409, 511, 514, 573, 685, 690, 758–759, 805
 Saupe order matrix, 109, 804
 Scalar coupling, 23–27, 55
 2Q and, 474–475
 active, 91, 282–284, 412, 456, 458, 473, 536

- apparent, 110, 660, 664–665, 669, 799
average, 402, 799
Bloch equations in, 23–27
chemical exchange and, 401–402
coherence transfer and, 280–283, 288, 338, 410, 492f, 520, 534, 601
CPMG and, 711, 714
dihedral angle and, 656, 799–802, 800f, 802f, 803f
dihedral angle restraints from, 455, 656, 798–803, 810
energy of spin states, 24
evolution in the product operator formalism, 84–85
Hamiltonian and, 61–65, 80, 378. *See also* Hamiltonian, scalar coupling
intramolecular dipolar relaxation and, 378–381
Karplus curve, 388, 424, 798–803, 800f, 802f, 803f
multiple quantum coherence, evolution of, 92, 536
multiplet, 24, 63, 76, 98, 201, 241–242, 271, 380, 401–402, 548, 571, 578, 684
passive, 91–92, 282, 655
 2Q in, 468
 2QF-COSY in, 443
 3Q in, 482
 COSY in, 412, 420
 COSY- β in, 426–427
 decoupled HSQC in, 555
 E.COSY in, 456–463
 HCCH-COSY in, 605
 PEP in, 561
relaxation and, 370, 387
resolved, 452
restraints derived from, 799–800
sign of, 459
spin echo and, 95
strong, 26–27, 62, 90, 107, 284–286, 292
time-dependent, 387–388
TOCSY and, 486
topology of, 490, 786
trans-hydrogen bond, 762, 806
in two-spin system, 25f, 62
weak, 26, 62, 64
- Scalar coupling constants, 24, 62, 84, 110, 401–402, 667t
 COSY and, 420–426
 HCCH-TOCSY and, 602
 heteronuclear, 617t
 homonuclear, 401, 504, 535, 538, 539, 541, 543, 583, 670, 703, 714, 726, 734, 798–802, 800f, 800t, 801t, 802f, 802t, 803f, 803f, 806–807
 linewidth and, 426
 Lorentzian lineshape and, 425
 measurement of, 420–426, 656–665
 side chains and, 656
 TOCSY and, 502
 trans-hydrogen bond, 797, 806
- Scalar coupling constants, measurements of
 absorptive and dispersive lineshapes from, 425
 COSY, 420–426
 COSY- β , 426–427
 E.COSY, 455–463
 heteronuclear E.COSY, 656
 heteronuclear-edited NMR, 590, 656
 HNCA-*J*, 656–660
 HNHA, 660–665
 nonlinear least squares, 426, 656
 quantitative *J*-correlation, 656
 TOCSY, 502
 TOCSY-HSQC, 590
- Scalar product, 33, 37
- Scalar relaxation, 370, 381, 387–388, 401, 406, 734
 of first kind, 387–388
 of second kind, 387–388
- Scaling interferogram, first point of, 155, 156f
- Schrödinger equation, 30, 41. *See also* Liouville von Neumann equation
 Hamiltonian and, 30
 time-independent, 34
 wavefunction and, 30
- SCUBA (Stimulated Cross-peaks Under Bleached Alphas), 426, 429
- Secondary chemical shift, 23, 804
- Secondary magnetic field, 22
- Secondary structure, 786, 790
 $^1\text{H}^{\text{N}}\text{--}^1\text{H}^{\alpha}$ scalar coupling for, 798–799

- Secondary structure (*Continued*)
 chemical shifts for, 405, 409, 783, 794, 804
 determination of, 806–808
 deuterium isotope shift, 733
 NOE characteristic for, 797, 806
 Secular, 355, 358, 360, 682
 Secular approximation, 358, 360–363, 372, 376, 380, 683
 SEDUCE-1, 170, 182f, 183f, 217
 Selection rule, 6, 25, 358, 449, 452–453, 473, 475, 482–483
 p-quantum, 482–483
 p-quantum filtration, 457
 Selective decoupling, 170, 217, 271, 549, 758
 Selective inversion recovery, 343, 346f
 Selective protonation. *See* Protonation
 Selective pulse, 172, 179–181, 217, 227, 228f, 229f, 230–232, 259, 525, 549, 607, 697
 Selectivity, phase cycle, 303–304
 Self-cancellation, 284, 414, 424, 425, 435, 476, 481, 510
 heteronuclear correlation and, 552
 Self-decoupling, relaxation by, 381, 401, 660, 799
 Semi-selective decoupling, 170, 549
 Sensitivity, 6, 98–99, 116–117, 119, 158.
 See also Noise; Signal-to-noise ratio; *individual NMR experiments*
 3D compared to 4D, 600–601
 constant-time, 544
 heteronuclear NMR, 534, 552
 recycle delay and, 257–258
 relaxation and, 534
 Sensitivity-enhancement. *See* PEP
 Separation of interactions, 278
 Separation of variables, 31
 Sequence alignment, 788, 790, 794
 Sequence specific assignment, 510, 788, 792, 796
 Sequential assignment, 515–516, 516f, 590, 614, 646, 653, 740, 785, 789, 791–793, 806
 Sequential NOE, 513f, 515, 533, 786f, 788–790, 790f, 807
 Shape factor, 220, 311
 Shaped pulse, 170, 171, 183f, 184f, 185f, 220, 228f, 230, 579, 645, 655, 696f, 698f, 717f
 Shielding tensor, 22, 102, 103
 Shift basis operator, 70, 73t, 75
 Shift operator, 70, 72, 73t, 75, 80–81, 90, 293, 316, 356
 Shifted laminar pulse (SLP). *See* Phase-modulated pulse
 Shim system, 117, 238–252
 coils and, 239–240, 240t, 250
 FID and, 241
 gradient, 242–252
 maps, 251f
 profile edge, 252
 protocol for, 243f
 pulse sequences for, 248f
 values, 251
 Shimming, 238–252
 Side chain conformation, 522, 656
 Side chain torsion angle, 656, 801–803, 803f, 803t
 Side chains, 19, 436, 667, 791
 ¹H and, 743
 2Q and, 478
 3Q and, 485, 486
 aliphatic, 641, 784f, 785
 CC(CO)NH-TOCSY and, 743
 deuteration and, 732, 740–743
 E.COSY and, 456
 nonaliphathic, 418, 784f, 785, 810
 R.COSY and, 433
 scalar coupling constants and, 656
 TOCSY and, 498, 499
 Sidebands. *See* Cycling sidebands
 Signal
 amplitude modulated, 316, 319, 322, 500, 559–560
 analog, 123, 127–129, 156
 audio frequency, 132
 continuous, 124
 cosine-modulated, 132, 315, 316, 320, 561, 562, 564, 771
 harmonic, 132, 144
 high-frequency, 126f, 132, 134f
 N-type, 316, 319, 322
 phase modulated, 316, 565
 P-type, 316, 319, 322
 sine-modulated, 132, 315, 316, 320–322, 326, 413, 561

- sinusoidal, 124, 132, 160, 254, 337, 411
- Signal averaging, 174, 255, 257, 258, 265, 274, 293, 310
- Signal-to-noise (S/N) ratio, 119, 123, 143, 144, 148, 149f, 150f, 158–159, 327–328. *See also* Noise; Sensitivity; *individual NMR experiments*
- 2QF-COSY and, 447, 448f
- COSY and, 414, 415, 448f
- deuteration and, 729–732, 741, 745
- ¹H NMR, in, 406–409
- pulsed field gradient, 313, 315, 322–323, 576
- heteronuclear NMR, 535
- linewidth and, 18
- matched filter and, 144
- multi-dimensional spectrum, in, 326
- probe, of, 119
- NOESY and, 506, 797
- PEP, 501, 560–566
- power ratio, 128
- preamplifier, of, 123
- rapid data acquisition and, 769, 770, 773, 775
- TOCSY and, 501
- ubiquitin 1D spectrum, in, 406
- Similarity theorem, 138
- Similarity transformation, 44, 48, 58
- Simulated annealing (SA), dynamical, 809
- Simultaneous eigenfunctions, 34–35
- Sinc function, 145, 146f, 182f
- Sinc pulse, 171, 182f
- Sinc-wiggles, 157
- Sine bell, 415f, 416, 417f, 426
 - 2QF-COSY and, 446
 - phase-shifted, 147
- Sine-modulated signal, 132, 315, 316, 320–322, 326, 413, 561
- Single element basis operator, 80–83, 99, 353, 450, 658
- Single quantum relaxation. *See* Spin-lattice relaxation; Spin-spin relaxation
- Single transition shift basis operator, 81, 83f, 100, 294, 460
- Single-quantum (SQ) coherence, 74, 82, 83f, 101f, 295
 - HSQC and, 535, 543–544, 572
 - operators and, 82, 88–90
- Single-quantum transitions, 76, 77f, 82, 83, 88, 202
- Size limitations, for NMR, 405, 533, 613
- Skewed unimolecular exchange, 399
- SL. *See* Spin lock
- Slices of 3D spectrum, 328–329
- Slow exchange, 401, 511, 522, 702, 754, 760. *See also* Chemical exchange
- Slow tumbling, 367, 391, 569
- SLP (Shifted laminar pulse). *See* Phase-modulated pulse
- S/N. *See* Signal-to-noise ratio
- Soft-hard-soft pulse sequence, 230
- Solomon equations, 338–346, 375, 503.
 - See also* Cross-relaxation; Dipolar relaxation
 - relaxation interference and, 386
- Solomon transition rate, 375
- Solvent exchange. *See* Amide proton exchange; Chemical exchange
- Solvent suppression, 117
 - 1-3-3-1, 224–227
 - binomial sequences, 224–227, 230
 - convolution difference low pass filter, 232–235
 - dynamic range and, 221, 223
 - flip-back, 231–232, 574–578, 575f, 685, 690–691
 - Hahn echo, 232, 263–267
 - heteronuclear NMR experiments and, 573–578
 - isotope exchange, 222–223
 - jump-return, 224–226, 227f, 233f, 235f, 261f
 - NOESY and, 503, 511–514
 - postacquisition, 232–234, 235f
 - presaturation, 181, 223–224, 233f, 409, 418, 428, 448, 473, 475, 499, 506, 511, 513f, 573
 - pulsed field gradient using, 227–232, 573, 574
 - saturation transfer during, 223, 226, 228, 232, 409, 573, 690
 - selective pulses in, 179–180, 227–232
 - spin lock purge pulse, 173, 227–230, 228f, 232, 573, 585, 767, 768
 - TOCSY-NOESY and, 528

- Solvent suppression (*Continued*)
 water flip-back, 231–232, 574–578, 575f, 685, 690–691
 WATERGATE, 229f, 230, 232, 574
Solvent, viscosity of, 21, 366, 408
Spatial variables, relaxation for, 353, 365, 368
Spatially dependent phase, 218, 220, 311, 576
Specific volume, 21
Spectral assignments, 782
Spectral density function, 355, 358, 363, 365–370, 367f
 autocorrelation for, 355
 cross-correlation for, 359, 369–370
 high-temperature limit in, 365
 internal motion for, 369
 lattice jump model for, 368
 Lipari-Szabo, 368–369
 rigid isotropic sphere, for, 366–367
Spectral editing, 546, 580
Spectral simplification, 437, 449, 452, 484
Spectral width, 124–125
 2Q experiment and, 473
 3Q NMR, 483
 ^{13}C , 550–551
Spectrometer. *See* NMR spectrometer
Spectrum, frequency domain, 18, 133–136, 773
Spherical harmonic functions, 106, 239, 240, 349, 365
 modified second-order, 350t
Spherical tensor operator, 103, 353
Spherical top, 366
Spin angular momentum. *See* Angular momentum
Spin decoupling. *See* Decoupling
Spin diffusion, 367, 511, 515, 528, 745, 759, 797, 798
 NOESY and, 506–508, 522
 ROESY and, 522
 regime, 367
Spin echo, 92–97, 265, 703, 711, 714
 chemical shifts and, 94
 difference, 763
 gradients and, 313, 315, 438
 in gradient shimming, 248
 Hamiltonian, 95
 homonuclear, 96
 multiple, 221
Spin flip, 338
Spin lock (SL), 197, 227–232, 228f, 254, 382, 518. *See also* Solvent suppression
 purge pulse, 173, 227–230, 228f, 232, 573, 585, 767, 768
 ROESY in, 520–524
 rotating-frame relaxation, in, 707–710
 trim pulse, 609
Spin operator, 46
Spin quantum number, 2
Spin system, 24, 28, 58. *See also* Three-spin system; Two-spin system
 AB, 26
 AX, 25f, 26
 categorization of, 787f
 density matrix formalism and, 29
 I_nS , 553, 555–556
Spin system assignments, 222, 405, 534, 613, 615t, 725, 739, 761, 782, 792
Spin tickling, 271
Spin-1/2 nuclei, 6, 27, 38, 55, 78, 370
Spin-lattice relaxation, 5, 12, 337, 338, 340, 378, 534, 805
 ^1H , 535
 axial peaks and, 309
 Bloch formulation, 12–14, 337
 CSA, 20f, 23, 349, 365, 366t, 370–371, 383–386, 680, 685
 dipolar, 346, 365, 366t, 370–378
 heteronuclear, 535
 inversion recovery, measuring using, 258–260, 343–346, 686, 734
 quadrupolar, 365, 366t, 370, 383–385, 680, 686, 693, 696, 700f
 scalar coupled spins, 378
 slow, 491
 Solomon equations, 338–346, 375, 386, 389
Spin-lattice relaxation rate constant, 5, 12, 337
Spin-spin coupling. *See* Scalar coupling
Spin-spin relaxation, 13, 337. *See also* Chemical exchange
 average, 364, 378–381
 Bloch formulation, 12–14, 337
 CSA, 20f, 23, 349–350, 365, 366t, 370–371, 383, 384t, 385t, 542, 569, 680, 685, 728, 729f, 739

- deuteration and, 726–728, 729f, 730f
dipolar, 19, 20f, 365, 366t, 370–371, 377–381, 739
heteronuclear, 542, 545, 557
linewidth, measuring from, 18–19, 406–408
quadrupolar, 2, 370, 383–385, 680, 686, 696
random-phase model for, 346–350
scalar, 387–388, 401, 406, 734
scalar coupled spins, 378–381
Spin-spin relaxation rate constant, 13, 337
Spin-state selective coherence transfer (S^3CT), 99, 100, 101f
Spontaneous emission, 334–335
SQ. *See* Single-quantum coherence
Stability, 116, 236, 660, 679
State function, 30–31, 34, 37. *See also* Eigenfunction; Wavefunction
States (hypercomplex) frequency discrimination, 315, 320, 322–325, 511, 550, 564, 576. *See also* TPPI, TPPI-States
Static magnetic field, 4, 7, 11, 116, 238, 246
Stationary state, 24, 34, 36, 55, 63, 75, 77, 98, 289, 336–337
Statistical mechanics, 40
Steady-state NOE difference experiment, 389–390, 690, 700
Steady-state NOE enhancement, 389
Stereo-array isotope labeling (SAIL), 727
Stereospecific assignments, 502, 522, 761, 798
Stimulated Cross-peaks Under Bleached Alphas (SCUBA), 426, 429
Stimulated emission, 334–335
Stochastic Brownian motion, 335, 365
Stochastic correlation function, 365. *See also* Spectral density function
 factoring, 367
 internal, 367
 overall, 367
 rigid isotropic sphere for, 366
Stochastic field, 335–336, 350–351, 365–367
Stochastic Hamiltonian, 345, 351–353, 358, 359, 370, 385, 679
Stochastic process, 336, 348, 355
Stoichiometry, 756
Stokes Law, 21
Straight-through coherence transfer, 632, 643, 743, 751
Strong coupling, 26, 62–64, 66, 90, 217, 285, 488, 611f
Strong coupling parameter, 62
Structural restraint. *See* Restraint
Structure determination, 806–811
 backbone conformation, 656
 back-calculation, 810
 CORMA, 810
 distance geometry, 808
 global fold, 749, 751, 786f, 807, 807f
 high resolution, 806–813
 restrained molecular dynamics, 808–809
 secondary structure, 529, 733, 751, 783, 786f, 790, 794, 804, 806–807, 813
 side chain conformation, 656
 simulated annealing, 809
 three-dimensional structure, 808–811, 813
Structure-activity relationship (SAR), 756
STUD, 211
Supercycle, 205, 206, 209, 211, 491, 493–494, 555, 557
Superoperator, 45, 351, 353, 356, 361, 363
Superposition state, 75, 90
SUSAN, 206, 208f, 209, 214, 215
Symmetrical reconversion, 684
Symmetry, 349, 361, 365, 373, 384, 387, 455, 456f, 588, 772, 783
Synchronous decoupling, 555–556
Synthesizer, frequency, 115f, 120, 189
- T**
- t_1 , 274
 T_1 . *See* Spin-lattice relaxation
 t_1 noise, 262, 453
 T_2 . *See* Spin-spin relaxation
 t_2 , 274
T5M4, 208–209
TALOS, 804
Temperature
 calibration, 236–237

- Temperature (*Continued*)
 - correlation time, effect on, 21
 - fluctuations, 310
- Tensor, 102–103, 104
 - chemical shift, 21–22, 102, 107
 - scalar coupling, 107
- Tensor operator
 - Cartesian, 102
 - CSA interaction and, 384t
 - dipolar interaction and, 372t
 - quadrupolar interaction and, 385t
 - spherical, 103–104, 353
- Tesla, 8
- Tetramethylsilane (TMS), 262
- Thermodynamic equilibrium, 55
- Three-dimensional (3D) shim map, 252
- Three-dimensional NMR. *See* 3D NMR
- Three-quantum (3Q) coherence, 442, 449–551, 481–482
- Three-spin single-quantum coherence, 450–451
- Three-spin system, 75–77, 77f, 91–92, 787f
 - 2Q and, 467–473, 471f
 - 2QF-COSY, and, 442–444
 - 3Q and, 481–483
 - 3QF-COSY, and, 449–451, 457–462
 - DCOSY, and 292
 - ROESY, in, 522
 - spin-diffusion in, 506–507
- Through-bond coherence transfer, 280
 - COSY-type, 281–284, 290–291
 - scalar coupling, by, 782
 - TOCSY-type, 284–289, 290–291
- Through-space coherence transfer, 280
 - NOE, by, 289–290
 - RDC, by, 291–292
- Tilt angle, 16
- Tilted rotating frame, 363–364, 364f, 381–383
- Time-dependent Hamiltonian, 30, 351
- Time-independent Hamiltonian, 31, 51, 351
- Time-proportional phase-incrementation (TPPI), 315, 321–322, 323t. *See also* States frequency discrimination; TPPI-States
 - folding and, 324, 325t, 550, 550f
 - initial delay for, 326
- Time-resolved fluorescence spectroscopy, 21
- Time-shifting theorem, 138, 152
- TMS (Tetramethylsilane), 262
- TOCSY (Total Correlation Spectroscopy), 284–289, 486–502. *See also* CC(CO)NH-TOCSY; Coherence transfer; HCCH-TOCSY; (H)CC(CO)NH-TOCSY; HSQC-TOCSY; Isotropic mixing; NOESY-TOCSY; TOCSY-HSQC
 - chemical exchange, 493
 - coherence transfer functions, 491, 492f, 495f
 - DCOSY, compared with, 292
 - disadvantages of, 498–499
 - dispersive lineshape in, 288, 490
 - experimental protocol for, 493–496
 - Hamiltonian for, 284, 287
 - heteronuclear edited NMR and, 582
 - isotropic mixing sequences and, 287–288, 487–493, 492f, 495f, 611f
 - Lorentzian-Gaussian transformation for, 496–497
 - mixing period and, 494–496, 495f, 497f
 - pre-TOCSY, 428–429, 428f, 515–517
 - processing of, 496–498
 - product operator analysis for, 284–289, 486–489
 - pulse sequence for, 488f, 489f
 - rapid acquisition, 500, 501f
 - relaxation in, 363–364, 496
 - relayed, 428–429, 428f, 515–517
 - ROE peaks in, 491, 493
 - sample purity, and, 409
 - sensitivity-enhanced, 500–502
 - stereospecific assignments, for, 502
 - ubiquitin, of, 480f, 497f, 501f
 - variants of, 428, 429, 499–502
 - zero-quantum peaks, 490
 - z-filtration, 288, 490
- TOCSY-HSQC, 589–591
 - pulse sequence for, 590f
 - resonance assignment and, 792
 - ubiquitin of, 591
- Torque, 7, 220

- Torsion-angle dynamics, 808
Torsion-angle, side-chain, 656, 801–803, 803f, 803t
Total Correlation Spectroscopy.
 See TOCSY
TPPI (Time-proportional phase-incrementation), 315, 321–322, 323t. *See also* States frequency discrimination; TPPI-States
TPPI-States, 322, 323t, 324–326, 325t, 655. *See also* States frequency discrimination; TPPI
Trace, 39
 basis operator, 79
 commutator, 40
 cyclic permutation, 40
 expectation value as, 39
Transferred NOE, 761
Transformation. *See also* Fourier transformation
 basis sets, between, 48
 coordinate, 108
 interaction frame, 352, 363
 product operator, of, 79, 84, 85f, 87f
 rotating frame, 10, 43–46, 50–51, 363–364
 similarity, 44–45, 48, 341
 spherical tensors, irreducible, of, 104
 tilted, 363–364, 364f, 517–519
 unitary, 44–45, 341
Transient NOE experiment, 289, 344, 346f, 390–391, 700, 805
Transient ROE experiment, 390, 391
Transition, conformational. *See* Chemical exchange
Transition rate constant, 345, 375
Transition
 allowed, 6, 25
 anticonnected, 460
 connected, 76, 456, 460
 double quantum, 75, 76f, 83f
 flip-flip, 75, 76f, 83f, 339, 346
 flip-flop, 75, 76f, 83f, 339, 346
 frequency, 6, 8
 magnetic dipole, 6
 multiple quantum, 76f
 non-connected, 460
 probability, spontaneous, 334–335
 progressively connected, 77, 460
 regressively connected, 77, 460
 single quantum, 25f, 75–77, 77f, 82
 stochastic field, induced by, 335–336
 three-spin system, in, 77f
 two-spin system, in, 25f, 83f
 unconnected, 426–427, 463
 Zeeman, 6
 zero-quantum, 75, 76f, 83f
Transmitter
 power attenuation, 253–254
 frequency, 9
 rf, 115f, 120–122
 triple-resonance experiments, in, 654
 tuning and matching, 237–238
Transverse relaxation. *See* Spin-spin relaxation
Transverse Relaxation Optimized Spectroscopy. *See* TROSY; Relaxation interference
Trim pulse. *See* Spin lock
Triple-quantum. *See* 3Q (Three-quantum)
Triple-resonance NMR, 613–655.
 See also individual NMR experiments
 3D and 4D, compared, 654–655
 experimental parameters, 654–655
 experiments, table of, 615t–617t
 nomenclature, 613
 out-and-back, 613
 out-and-stay, 632
 straight-through, 632
TROSY (Transverse Relaxation Optimized Spectroscopy), 387, 535, 552–572, 574–578. *See also* Relaxation interference
 calbindin D_{28k}, of, 572
 chemical shift mapping, for, 756
 coupling-enhanced TROSY (CE-TROSY), 672–673, 673f
 CPMG and, 720, 721f
 cross saturation, for, 758f
 energy level representation for, 559f
 Hahn echo and, 718, 719f
 methyl groups, for, 749, 770
 phase cycling and, 122, 555, 567
 product operator analysis for, 566–568
 pulse sequence for, 554f, 575f, 673f, 758f

- TROSY (*Continued*)
 pulsed field gradients in, 574–578, 577f
 RDC, for measuring, 672
 relaxation and, 386–387, 568–572,
 719–720
 ubiquitin, of, 570f
 water flip-back in, 574, 575f
 TROSY-CPMG, 720, 721f
 TROSY-HNCA, gradient-enhanced,
 620f, 628
 Truncation artifacts, 143–148, 146f, 157,
 265
 Tuning and matching, 237–238, 238f
 Two-dimensional NMR. *See* 2D NMR
 Two-quantum. *See* 2Q (Double-
 quantum)
 Two-spin order, 314, 519, 574, 699, 706,
 720
 Two-site exchange, 392, 400f, 755f
 longitudinal magnetization, 394–397
 rate matrix, 393, 395, 397
 transverse magnetization, 395,
 397–399, 708, 711 755f
 Two-spin system, 25f
 energy, 24–26, 63, 371–372
 Hamiltonian for, 62, 64
 isolated, 345, 507
 rotation matrices, 66–67
 2Q and, 466, 467f
 Cartesian basis, 70
 direct product spaces in, 59
 one-pulse experiment for, 68–70
 product operators for, 71t, 73t
 dipolar relaxation rate constants
 for, 377t
 scalar coupling in, 25f, 62, 64
 transitions for, 25f, 339f, 371f
 single element basis, 81–82, 83f
 wavefunctions, 24, 59, 62
 Tycko 5-step phase cycle, 208–209, 211

U

Ubiquitin

- chemical shifts in, 783f
- correlation time of, 21
- denatured, 407f, 408
- description, vi, ix
- linewidth for, 408
- order parameters for, 812

- RDCs for, 669f, 812f
 - relaxation decay for, 693f, 710f
 - relaxation dispersion in, 710f
 - relaxation rate constants for, 694f,
 695f, 700f, 705f, 717f
 - sample preparation, ix
 - scalar coupling constants for, 667t,
 800f
 - structure, NMR-derived, 807f, 811f
- Ubiquitin, NMR spectra of
 heteronuclear
 HCCH-COSY, 609f
 HCCH-TOCSY, 612f
 HMQC, 544f
 HNHA, 666f
 HSQC, 544f, 551f, 570f, 581f, 582f,
 669f
 HSQC-NOESY-HSQC, 747f, 752f
 NOESY-HSQC, 587f, 589f
 TOCSY-HSQC, 591f
 TROSY, 570f
- homonuclear
 2Q, 467f, 477f–480f
 2QF-COSY, 445f, 446f, 449f, 450f,
 453f, 454f, 462f, 480f
 3Q, 484f, 487f
 3QF-COSY, 453f–455f, 462f
 COSY, 415f, 417f, 419f–422f, 427f,
 449f
 COSY-35, 427f, 464f
 DR.COSY, 436f
 E.COSY of, 462f, 464f
 JR.NOESY, 513f
 NOESY, 509f, 513f, 514f, 516f
 pre-TOCSY COSY, 428f
 R.COSY, 434f
 relayed NOESY, 516f
 ROESY, 523f
 TOCSY, 480f, 497f, 501f
- 1D
 excitation sculpting, 233f
 Hahn echo, 267f, 407f, 783f
 inversion recovery, 260f
 jump-return, 233f, 261f
 1D, denatured, 407f
 one-pulse acquire, 150f, 233f, 235f,
 267f, 272f
 spin-echo, 261f, 267f, 407f, 783f
 triple-resonance
 CBCA(CO)NH, 646f

- CBCANH, 650f, 774f
 - CC(CO)NH-TOCSY, 744f
 - HACA(CO)NH, 774f
 - HACANH, 774f
 - H(CA)NH, 636f
 - HNCA, 624f
 - HNCACB, 654f
 - HNCA-*J*, 661f
 - HN(CO)CA, 632f
 - Unconnected transition, 426–427, 463
 - Unit vector, 5, 9–10, 54, 365
 - Unitary matrix, 341
 - Unitary operator, 44
 - Unitary transformation, 44, 45
 - Unlike spins, 360–363, 714
- V**
- van der Waals, 798, 809
 - Variable target function distance geometry, 808
 - Vector space, 32, 33, 38, 60, 79
 - Vicinal proton, 796–798
 - Viscosity, 21, 366, 408
 - Voltage, 124, 237, 253–254
 - Voltage standing-wave ratio (VSWR), 237
 - VSWR (Voltage standing-wave ratio), 237
- W**
- WALTZ, 205–206, 207f, 209, 214f, 217
 - synchronous, 555, 627, 657
 - Hahn echo and, 703
 - HSQC and, 555
 - Isotropic mixing sequence, as, 490–491, 492f
 - Saturation by, 690
 - WALTZ-4, 205, 207f
 - WALTZ-8, 205, 207f
 - WALTZ-16, 205–206, 207f, 214f, 490–491
 - Water, bound, 512–513
 - Water, chemical shift, 237
 - Water flip-back, 231–232, 574–578, 575f
 - saturation transfer and, 690–691
 - Water radiation damping. *See* Radiation damping
 - Water stripe, 416, 477
 - Water suppression. *See* Solvent suppression
 - WATERGATE, 229f, 230, 232, 574, 671f
 - Wavefunction, 24, 30–33, 35, 37–38, 40, 47–48, 59, 62–64, 74
 - Weak scalar coupling, 26, 62, 64, 80, 90, 284, 538
 - weight factor, E.COSY, 457
 - WHATIF, 809–810
 - Wigner rotation matrices, 104–107, 105t
 - Wigner-Eckart theorem, 3
 - Window function. *See* Apodization function
 - Windowing. *See* Apodization
 - WURST, 198–200, 200f
 - isotropic mixing, 491–493, 492f
 - spin decoupling, 211–212, 213f, 216f, 758
- Z**
- Zeeman Hamiltonian, 50, 55–57, 355
 - Zeeman levels, 4, 6
 - Zeeman transition, 8, 12
 - Zero-filling, 142–143, 600
 - Zero-order phase correction, 152, 154f, 322, 326, 416, 563–564
 - Zero-padding, 142–143, 600
 - Zero-quantum (ZQ) coherence, 74, 75, 76f, 82, 83f, 90–92, 314
 - chemical exchange and, 715
 - relaxation of, 540
 - scalar coupling and, 92
 - two-pulse segment for, 306f
 - Zero-quantum peak, 490, 505–506, 508–509
 - suppression of, 314, 508–509
 - Zero-quantum splitting, 92
 - Zero-quantum (ZQ) transition. *See* Flip-flop transition
 - Z-filtration, 288, 490, 496
 - Z-gradient, 218–220. *See also* Pulsed field gradient
 - ZQ. *See* Zero-quantum coherence
 - Z-rotation, 300
 - ZZ-exchange, 702, 706–707
 - pulse sequence for, 707f