

LIST OF FIGURES

- Figure 1.1. Angular momentum
- Figure 1.2. Orientations of magnetic fields in the rotating reference frame
- Figure 1.3. Effect of an applied rf field on magnetization
- Figure 1.4. Effects of on-resonance pulses
- Figure 1.5. One-pulse NMR experiment
- Figure 1.6. Resonance linewidths
- Figure 1.7. Energy levels for AX spin system
- Figure 2.1. Multiple quantum transitions for an *IS* spin system
- Figure 2.2. Spin states and spectrum for a three-spin *IRS* system
- Figure 2.3. Geometrical representations of rotations in an operator space
- Figure 2.4. Single-transition basis operators for an *IS* spin system
- Figure 2.5. Operator rotations
- Figure 2.6. Transformations of product operators
- Figure 2.7. Energy-level diagram for S³CT pulse sequence
- Figure 2.8. Coordinate transformations by Euler angles
- Figure 3.1. Block diagram of an NMR spectrometer
- Figure 3.2. Diagram of a superconducting magnet
- Figure 3.3. Probe assembly
- Figure 3.4. Nyquist theorem
- Figure 3.5. Experimental scheme for quadrature detection

- Figure 3.6. Illustration of quadrature detection
- Figure 3.7. Adsorptive and dispersive Lorentzian lineshapes
- Figure 3.8. Window functions for apodization
- Figure 3.9. Digital resolution enhancement
- Figure 3.10. Digital resolution enhancement of ubiquitin ^1H NMR spectrum
- Figure 3.11. Phase dependence of lineshapes
- Figure 3.12. Phase corrections
- Figure 3.13. Baseline distortions from phase corrections
- Figure 3.14. Baseline distortions from a corrupt FID
- Figure 3.15. Linear prediction using the HSVD algorithm
- Figure 3.16. Linear prediction of an HSQC spectrum
- Figure 3.17. Maximum entropy reconstruction of an HSQC spectrum
- Figure 3.18. Off-resonance effects for 90° and 180° pulses
- Figure 3.19. Resonance lineshapes for off-resonance 90° pulses
- Figure 3.20. Excitation profiles for 180° pulses
- Figure 3.21. Selective pulse shapes
- Figure 3.22. Selective 90° pulses
- Figure 3.23. Selective 180° pulses
- Figure 3.24. Phase-modulated pulse
- Figure 3.25. Vector representations of phase-modulated pulses
- Figure 3.26. Magnetization trajectories during an adiabatic pulse
- Figure 3.27. Pulse shape and excitation profile for a WURST-20 adiabatic sweep
- Figure 3.28. Scalar coupling scaling factor for WALTZ decoupling sequences
- Figure 3.29. Scalar coupling scaling factor for GARP-1 and SUSAN-1 decoupling sequences
- Figure 3.30. Scalar coupling scaling factor for MPF-7 and MPF-9 decoupling sequences
- Figure 3.31. Scalar coupling scaling factors for WURST-40 decoupling sequence
- Figure 3.32. Cycling sidebands for WALTZ-16, GARP-1, and SUSAN-1 decoupling sequences
- Figure 3.33. Cycling sidebands for MPF-7 and MPF-9 decoupling sequences
- Figure 3.34. Cycling sidebands for WURST decoupling sequences
- Figure 3.35. Dephasing of transverse magnetization by pulsed field gradients
- Figure 3.36. Intrinsic backbone H^{N} exchange rates
- Figure 3.37. Binomial excitation profiles
- Figure 3.38. Water suppression using spin lock purge pulses

- Figure 3.39. Water suppression using field gradient pulses
Figure 3.40. Excitation profiles for water-selective pulses
Figure 3.41. Experimental examples of water suppression techniques
Figure 3.42. Post-acquisition solvent suppression by low-pass filtration
Figure 3.43. Tuning circuits
Figure 3.44. Shimming protocol
Figure 3.45. Gradient image of transmitter/receiver coil volume
Figure 3.46. Pulse sequences for z -axis gradient shimming
Figure 3.47. Typical z -axis shim maps
Figure 3.48. Pulse sequences for indirect calibration of heteronuclear pulse lengths
Figure 3.49. Ernst angle
Figure 3.50. ^1H inversion recovery spectra of ubiquitin
Figure 3.51. Pulse sequence for 1D jump-return Hahn echo experiment
Figure 3.52. Estimating ^1H R_2 relaxation rate constants
Figure 3.53. Hahn-echo ^1H NMR spectra
Figure 3.54. Pulse-acquire and Hahn-echo ^1H NMR spectra of ubiquitin
Figure 4.1. ^1H NMR spectra of a hexapeptide and ubiquitin
Figure 4.2. General scheme for two-dimensional NMR spectroscopy
Figure 4.3. Dependence of the NMR signal on the t_1 evolution period
Figure 4.4. Schematic two-dimensional NMR spectrum for two spins
Figure 4.5. Coherence transfer pathway for a double quantum experiment
Figure 4.6. Pulse phase
Figure 4.7. Receiver phase
Figure 4.8. Pulse and receiver phase
Figure 4.9. Selection of double quantum coherence
Figure 4.10. Vectorial picture of phase cycling
Figure 4.11. Two-pulse segment used to generate zero quantum coherence
Figure 4.12. EXORCYCLE
Figure 4.13. Selection of a coherence transfer pathway by pulsed field gradients
Figure 4.14. Frequency discrimination in the F_1 dimension
Figure 4.15. Comparison of absorptive and phase twisted lineshapes
Figure 4.16. Folding and aliasing in the F_1 dimension
Figure 4.17. Schematic generation of a three-dimensional NMR experiment
Figure 4.18. Development of a three-dimensional data set from a two-dimensional data set

- Figure 4.19. Schematic representation of a four-dimensional NMR experiment
- Figure 4.20. Visualization of a four-dimensional experiment
- Figure 5.1. Transitions and associated rate constants for a two-spin system
- Figure 5.2. Time dependence of matrix coefficients for longitudinal relaxation
- Figure 5.3. Magnetization decays for inversion recovery experiments
- Figure 5.4. Redfield kite
- Figure 5.5. Relative orientations of laboratory and tilted reference frames
- Figure 5.6. Spectral density functions for an isotropic rotor
- Figure 5.7. Transitions and associated eigenfrequencies for a two-spin system
- Figure 5.8. Relaxation rate constants for ^1H - ^{15}N dipolar spin system
- Figure 5.9. Population transfer due to chemical exchange
- Figure 5.10. Chemical exchange for a two-site system
- Figure 6.1. One-dimensional ^1H spectrum of ubiquitin
- Figure 6.2. Pulse sequence and coherence level diagram for the COSY experiment
- Figure 6.3. Variation in the peak height of an anti-phase absorptive doublet
- Figure 6.4. Sections of the $^1\text{H}^{\text{N}}$ - $^1\text{H}^{\alpha}$ region of COSY spectra for ubiquitin
- Figure 6.5. Effect of t_1 window functions on COSY spectra
- Figure 6.6. Five fingerprint regions of the COSY spectrum of ubiquitin
- Figure 6.7. Leucine, valine and isoleucine methyl fingerprint region of the COSY spectrum of ubiquitin
- Figure 6.8. Alanine and threonine methyl fingerprint region of the COSY spectrum of ubiquitin
- Figure 6.9. $^1\text{H}^{\text{N}}$ - $^1\text{H}^{\alpha}$ fingerprint region of the COSY spectrum of ubiquitin
- Figure 6.10. Linewidth dependence of a Lorentzian anti-phase absorptive doublet lineshape
- Figure 6.11. Linewidth dependence of peak height of a Lorentzian anti-phase absorptive doublet
- Figure 6.12. Linewidth dependence of peak separation for anti-phase absorptive Lorentzian doublet
- Figure 6.13. Measurement of the peak separation for absorptive and dispersive anti-phase Lorentzian doublet

- Figure 6.14. Cross-peaks for Gly10 in the COSY-35 spectrum of ubiquitin
- Figure 6.15. Two sections of the pre-TOCSY COSY spectrum of ubiquitin
- Figure 6.16. Pulse sequence and coherence level diagram for the R.COSY experiment
- Figure 6.17. R.COSY spectrum of ubiquitin
- Figure 6.18. Pulse sequence and coherence level diagram for the DR.COSY experiment
- Figure 6.19. DR.COSY spectrum of ubiquitin
- Figure 6.20. Pulse sequence and coherence level diagram for 2QF-COSY experiment
- Figure 6.21. Pulse sequence and coherence level diagram for gradient-enhanced 2QF-COSY experiment
- Figure 6.22. 2QF-COSY spectrum of ubiquitin acquired with a short recycle delay
- Figure 6.23. Cross-peaks obscured by “double diagonal” artifacts
- Figure 6.24. Comparison of the S/N ratio in COSY and 2QF-COSY spectra
- Figure 6.25. Comparison of cross-peaks close to the diagonal in COSY and 2QF-COSY
- Figure 6.26. Gradient-enhanced 2QF-COSY spectrum of ubiquitin
- Figure 6.27. Methyl fingerprint region in 2QF and 3QF COSY spectra of ubiquitin
- Figure 6.28. $^1\text{H}^\alpha$ - $^1\text{H}^\beta$ cross-peaks in 2QF-COSY and 3QF-COSY spectra of ubiquitin
- Figure 6.29. $^1\text{H}^\alpha$ - $^1\text{H}^{\beta'}$ and $^1\text{H}^\alpha$ - $^1\text{H}^{\beta''}$ cross-peaks of Glu24 in the 3QF-COSY spectrum of ubiquitin
- Figure 6.30. Schematic view of partially overlapped cross-peaks in 2QF- and 3QF-COSY spectra
- Figure 6.31. Schematic view of cross-peaks for a three-spin system in 2QF-COSY, 3QF-COSY, and E.COSY spectra
- Figure 6.32. Section of the E.COSY spectrum of ubiquitin
- Figure 6.33. Comparison of the COSY-35 and E.COSY spectra of ubiquitin
- Figure 6.34. Pulse sequence and coherence level diagram for 2Q and 3Q experiments
- Figure 6.35. Example of the appearance of 2Q cross-peaks arising from two-spin systems
- Figure 6.36. Lineshapes of single and double anti-phase Lorentzian lines

- Figure 6.37. Contributions of initial I_1 magnetization to peaks observed in the 2Q spectrum of a three-spin system
- Figure 6.38. Schematic representation of the peaks expected in the 2Q spectrum of a $^1\text{H}^{\text{N}}\text{-}^1\text{H}^{\alpha}\text{-}^1\text{H}^{\beta 2}$ spin system
- Figure 6.39. $^1\text{H}^{\text{N}}\text{-}^1\text{H}^{\alpha}$ region of the 2Q spectrum of ubiquitin
- Figure 6.40. 2Q spectrum of ubiquitin showing remote glycine peaks at $F_1 = ^1\text{H}^{\alpha'} + ^1\text{H}^{\alpha''}$
- Figure 6.41. Remote and direct peaks for $\text{C}^{\alpha}\text{H}\text{-}\text{C}^{\beta}\text{H}_2$ fragments in the 2Q spectrum of ubiquitin
- Figure 6.42. Cross-peaks between $^1\text{H}^{\alpha}$ of Glu64 and its side-chain resonances in 2Q, TOCSY, and 2QF-COSY spectra of ubiquitin
- Figure 6.43. Spectral simplification achieved for three coupled spins in a 3Q spectrum of ubiquitin
- Figure 6.44. Section of the 3Q spectrum of ubiquitin showing aromatic side-chain resonances
- Figure 6.45. Pulse sequence and coherence level diagram for the TOCSY experiment
- Figure 6.46. Pulse sequence and coherence level diagram for a TOCSY experiment using excitation sculpting
- Figure 6.47. Efficiency of coherence transfer between two scalar coupled spins under the influence of different Hartmann-Hahn mixing schemes
- Figure 6.48. TOCSY cross-peak intensity as a function of isotropic mixing time
- Figure 6.49. TOCSY spectra of ubiquitin acquired with different mixing times
- Figure 6.50. Rapid acquisition of TOCSY spectra for ubiquitin
- Figure 6.51. Pulse sequence and coherence level diagram for the NOESY experiment
- Figure 6.52. Pulse sequence and coherence level diagram for the NOESY experiment using excitation sculpting
- Figure 6.53. NOESY spectra of ubiquitin acquired with different mixing times
- Figure 6.54. Pulse sequence and coherence level diagram for the JR-NOESY experiment
- Figure 6.55. NOESY spectra of ubiquitin with presaturation and JR water suppression
- Figure 6.56. NOESY spectra of ubiquitin with excitation-sculpting water suppression
- Figure 6.57. Pulse sequence and coherence level diagram for the relayed NOESY experiment

- Figure 6.58. Comparison of NOESY and relayed NOESY spectra of ubiquitin
- Figure 6.59. Pulse sequences and coherence level diagram for ROESY experiments
- Figure 6.60. ROESY spectra of ubiquitin
- Figure 6.61. Pulse sequence and coherence level diagram for a ROESY experiment using excitation sculpting
- Figure 6.62. Pulse sequence and coherence level diagram for a homonuclear 3D NOESY-TOCSY experiment
- Figure 7.1. Pulse sequences for HMQC, HSQC and constant-time HSQC experiments
- Figure 7.2. ^1H - ^{15}N HMQC, HSQC and constant-time HSQC spectra of ^{15}N -labeled ubiquitin
- Figure 7.3. Interferograms and lineshapes for ^1H - ^{15}N HMQC, HSQC and constant-time HSQC spectra
- Figure 7.4. Schematic illustration of halving the F_1 spectral width in a ^1H - ^{13}C heteronuclear correlation experiment
- Figure 7.5. ^1H - ^{13}C HSQC spectra of uniformly $^{15}\text{N}/^{13}\text{C}$ -labeled ubiquitin
- Figure 7.6. Pulse sequences for decoupled HSQC, sensitivity-enhanced PEP HSQC, and TROSY experiments
- Figure 7.7. Refocused INEPT coherence transfer functions for I_nS spin systems
- Figure 7.8. Energy-level representation of reverse polarization transfers for decoupled HSQC, TROSY and PEP-HSQC experiments
- Figure 7.9. ^1H - ^{15}N decoupled HSQC, PEP HSQC, and TROSY spectra of ^{15}N -labeled ubiquitin
- Figure 7.10. ^1H - ^{15}N multiplet structure for calbindin D_{28k}
- Figure 7.11. Pulse sequences for fast HSQC and TROSY experiments incorporating water-flipback solvent suppression
- Figure 7.12. Pulse sequences for PFG-PEP-HSQC and PFG-TROSY experiments
- Figure 7.13. Pulse sequences for the ^1H - ^{13}C constant-time HSQC experiment
- Figure 7.14. ^1H - ^{13}C HSQC and constant-time HSQC spectra of $^{15}\text{N}/^{13}\text{C}$ -labeled ubiquitin
- Figure 7.15. Expansions of the ^1H - ^{13}C HSQC and constant-time HSQC spectra of $^{15}\text{N}/^{13}\text{C}$ -labeled ubiquitin
- Figure 7.16. Pulse sequence for the 3D ^1H - ^{15}N NOESY-HSQC experiment

- Figure 7.17. Schematic illustration showing the relationship between a 3D heteronuclear-edited spectrum and 2D homonuclear and heteronuclear correlation spectra
- Figure 7.18. Selected $F_1(^1\text{H}^\alpha)$ - $F_3(^1\text{H}^\text{N})$ regions from a 3D ^1H - ^{15}N NOESY-HSQC spectrum of ^{15}N -labeled ubiquitin
- Figure 7.19. Selected $F_1(^1\text{H})$ - $F_3(^1\text{H}^\alpha)$ regions from a 3D ^1H - ^{13}C NOESY-HSQC spectrum of ^{13}C -labeled ubiquitin
- Figure 7.20. Pulse sequence for the 3D ^1H - ^{15}N TOCSY-HSQC experiment
- Figure 7.21. ^1H - ^{15}N TOCSY-HSQC spectrum of ^{15}N -labeled ubiquitin
- Figure 7.22. Pulse sequence for the HSQC-NOESY experiment
- Figure 7.23. Pulse sequence for the 3D ^1H - ^{15}N HMQC-NOESY-HMQC experiment
- Figure 7.24. Pulse sequence for the 4D $^{13}\text{C}/^{15}\text{N}$ HMQC-NOESY-HMQC experiment
- Figure 7.25. Pulse sequence for the 4D $^{13}\text{C}/^{13}\text{C}$ HMQC-NOESY-HMQC experiment
- Figure 7.26. Pulse sequences for 3D HCCH-COSY experiments
- Figure 7.27. Selected $F_1(^1\text{H})$ - $F_3(^1\text{H})$ slice from a constant-time HCCH-COSY spectrum of ^{13}C -labeled ubiquitin
- Figure 7.28. Pulse sequence for the 3D HCCH-TOCSY experiment
- Figure 7.29. Isotropic mixing for ^{13}C spins in isoleucine residues
- Figure 7.30. Selected regions from $F_2(^{13}\text{C})$ slices of the 3D HCCH-TOCSY spectrum of ^{13}C -labeled ubiquitin
- Figure 7.31. Pulse sequences for the HNCA experiment
- Figure 7.32. Plots of the HNCA coherence transfer functions, $\Gamma_k(T)$
- Figure 7.33. Plots of the HNCA coherence transfer functions, $\Gamma_k(\delta)\cos(\pi J_{\text{NH}}\delta)$
- Figure 7.34. Selected $F_2(^{13}\text{C}^\alpha)$ - $F_3(^1\text{H}^\text{N})$ slice from an HNCA spectrum of ubiquitin
- Figure 7.35. Pulse sequence for the constant-time HN(CO)CA experiment
- Figure 7.36. Selected $F_2(^{13}\text{C}^\alpha)$ - $F_3(^1\text{H}^\text{N})$ slice from a HN(CO)CA spectrum of ubiquitin
- Figure 7.37. Pulse sequence for the constant-time H(CA)NH experiment
- Figure 7.38. Plots of the H(CA)NH coherence transfer function, $\Gamma_\alpha(2\delta_2)$
- Figure 7.39. Selected $F_1(^1\text{H}^\alpha)$ - $F_3(^1\text{H}^\text{N})$ slice from an H(CA)NH spectrum of ubiquitin
- Figure 7.40. Pulse sequence for the constant-time HNCO experiment

- Figure 7.41. Pulse sequence for the constant-time HN(CA)CO experiment
- Figure 7.42. Pulse sequence for the CBCA(CO)NH experiment
- Figure 7.43. CBCA(CO)NH spectrum of ubiquitin
- Figure 7.44. Pulse sequence for the CBCANH experiment
- Figure 7.45. Plots of the CBCANH coherence transfer functions
- Figure 7.46. CBCANH spectrum of ubiquitin
- Figure 7.47. Pulse sequence for the HNCACB experiment
- Figure 7.48. HNCACB spectrum of ubiquitin
- Figure 7.49. Pulse sequence for the HNCA-*J* experiment
- Figure 7.50. Selected $F_1(^{13}\text{C}^\alpha)$ - $F_3(^1\text{H}^\text{N})$ slice from a HNCA-*J* spectrum of ubiquitin
- Figure 7.51. Pulse sequence for the HNHA experiment
- Figure 7.52. Effects of relaxation on the HNHA experiment
- Figure 7.53. HNHA spectrum of ubiquitin
- Figure 7.54. Residual dipole couplings in ubiquitin
- Figure 7.55. Pulse sequence for the IPAP ^{15}N - ^1H HSQC experiment
- Figure 7.56. Pulse sequence for the CE-TROSY experiment incorporating *J*-scaling
- Figure 8.1. Pulse sequences for ^{15}N R_1 , R_2 , $R_{1\rho}$ and *NOE* relaxation measurements
- Figure 8.2. ^{15}N relaxation data for ubiquitin
- Figure 8.3. ^{15}N relaxation rate constants for ubiquitin
- Figure 8.4. ^{15}N CSA/ ^1H - ^{15}N dipolar relaxation interference
- Figure 8.5. Pulse sequences for ^2H R_1 and $R_{1\rho}$ relaxation measurements in CH_2D spin systems
- Figure 8.6. Pulse sequences for ^2H quadrupolar order and antiphase coherence relaxation measurements in CH_2D spin systems
- Figure 8.7. ^2H relaxation rate constants for CH_2D methyl groups in ubiquitin
- Figure 8.8. Pulse sequences for ^{13}CO R_1 and $\{^{13}\text{C}^\alpha\}$ - ^{13}CO *NOE* relaxation measurements
- Figure 8.9. Pulse sequence for the in-phase Hahn-echo experiment
- Figure 8.10. Values of ^{15}N R_{ex} for ubiquitin
- Figure 8.11. Pulse sequence for the X nucleus ZZ-exchange experiment
- Figure 8.12. Pulse sequences for ^{15}N $R_{1\rho}$ relaxation measurements
- Figure 8.13. ^{15}N $R_{1\rho}$ relaxation dispersion for residue Asn 25 in ubiquitin
- Figure 8.14. Pulse sequences for relaxation-compensated CPMG experiments
- Figure 8.15. Pulse sequence for the $^1\text{H}^\text{N}$ CPMG experiment in ^{15}N - ^2H labeled proteins

- Figure 8.16. Multiple quantum relaxation in ubiquitin
- Figure 8.17. TROSY-based methods for detecting ^{15}N R_{ex} in large, deuterated proteins
- Figure 8.18. Pulse sequence for the ^{15}N TROSY-CPMG experiment
- Figure 9.1. T_2 relaxation times for fully protonated and perdeuterated proteins
- Figure 9.2. Transverse and longitudinal relaxation times as a function of deuteration level in proteins
- Figure 9.3. Pulse sequence for the 3D constant-time HNCA experiment
- Figure 9.4. Pulse sequence for the 3D HN(CA)CB experiment
- Figure 9.5. Pulse sequence for the 3D gradient-enhanced, sensitivity-enhanced CC(CO)NH-TOCSY experiment
- Figure 9.6. 3D CC(CO)NH-TOCSY spectrum of perdeuterated $^2\text{H}/^{13}\text{C}/^{15}\text{N}$ -labeled calbindin $\text{D}_{28\text{K}}$
- Figure 9.7. Pulse sequence for the gradient-enhanced, sensitivity-enhanced 4D $^{15}\text{N}/^{15}\text{N}$ -separated NOESY experiment
- Figure 9.8. 4D $^{15}\text{N}/^{15}\text{N}$ -separated NOESY spectrum of ILV-protonated, otherwise perdeuterated, calbindin $\text{D}_{28\text{K}}$
- Figure 9.9. Pulse sequence for the gradient-enhanced, sensitivity-enhanced 4D $^{13}\text{C}/^{15}\text{N}$ separated NOESY experiment
- Figure 9.10. 3D $^{13}\text{C}/^{13}\text{C}$ -separated NOESY, 4D $^{13}\text{C}/^{15}\text{N}$ -separated NOESY, and 4D $^{15}\text{N}/^{15}\text{N}$ -separated NOESY spectra of ILV-protonated, otherwise perdeuterated, calbindin $\text{D}_{28\text{K}}$
- Figure 9.11. Ligand-induced contributions to chemical exchange linebroadening
- Figure 9.12. TROSY pulse sequence for using cross-saturation to identify protein-ligand interfaces
- Figure 9.13. Basic pulse sequence for the ^{13}C double-half-filtered NOESY experiment
- Figure 9.14. Projection reconstruction NMR spectra of $^{13}\text{C}/^{15}\text{N}$ -labeled calbindin $\text{D}_{28\text{K}}$
- Figure 10.1. ^1H chemical shift ranges for ubiquitin
- Figure 10.2. Aliphatic and non-aliphatic side chain ^1H chemical shifts
- Figure 10.3. Assignment methods for unlabeled proteins.
- Figure 10.4. Categorization of spin-systems using chemical shifts and spin topology
- Figure 10.5. Backbone-backbone NOEs for sequence-specific ^1H resonance assignments
- Figure 10.6. Summary of sequential NOEs observed for a portion of ubiquitin
- Figure 10.7. Aliphatic and aromatic ^{13}C chemical shifts

- Figure 10.8. Karplus curves describing the variation of $^3J_{COCO}$, $^3J_{COH^\alpha}$, $^3J_{COC^\beta}$, $^3J_{H^NCO}$, $^3J_{H^NH^\alpha}$, and $^3J_{H^NC^\beta}$ with backbone dihedral angle ϕ
- Figure 10.9. Karplus curve describing the variation of $^3J_{H^\alpha N}$ with backbone dihedral angle ψ
- Figure 10.10. Karplus curves describing the variation of $^3J_{H^\alpha H^{\beta 2}}$ and $^3J_{H^\alpha H^{\beta 3}}$
- Figure 10.11. Definition of the global fold using long-range NOEs
- Figure 10.12. NMR-derived structure of ubiquitin
- Figure 10.13. Residual dipolar coupling constants for ubiquitin
- Figure 10.14. Backbone ^{15}N generalized order parameters for ubiquitin